

SOLE INVENTOR

"EXPRESS MAIL" Mailing Label No.  
EL542916975US

Date of Deposit: October 24, 2000.

I hereby certify that this paper (or fee),  
postage prepaid, is being deposited with the  
United States Postal Service "EXPRESS MAIL  
POST OFFICE TO ADDRESSEE" service under  
37 CFR §1.10 on the date indicated above  
and is addressed to: Assistant Commissioner  
for Patents, Washington, D.C. 20231

  
Sonji Shivers

APPLICATION FOR  
UNITED STATES LETTERS PATENT

S P E C I F I C A T I O N

-----

Attorney's Docket No. 11016US05 / 100-236.P2.C2

TO ALL WHOM IT MAY CONCERN:

Be it known that I, LEWIS H. LAMBERT, Jr., a citizen of the United States, residing at 45928 Omega Drive, Fremont, California 94539, a citizen of the United States, have invented new and useful "IMPROVED THERAPEUTIC COMPOSITIONS COMPRISING BACTERICIDAL/PERMEABILITY-INCREASING (BPI) PROTEIN PRODUCTS" of which the following is a specification.

004201" 60796960

**IMPROVED THERAPEUTIC COMPOSITIONS COMPRISING  
BACTERICIDAL/PERMEABILITY-INCREASING  
(BPI) PROTEIN PRODUCTS**

This is a continuation-in-part of U.S. Application Serial No.  
5 08/530,599 filed September 19, 1995, which is in turn a continuation-in-part of  
U.S. Application Serial No. 08/372,104 filed January 13, 1995, all of which are  
incorporated herein by reference.

**BACKGROUND OF THE INVENTION**

The present invention relates generally to improved therapeutic  
10 compositions and treatment methods utilizing poloxamer (polyoxypropylene-  
polyoxyethylene block copolymer) surfactants for enhancing the activity of  
bactericidal/permeability-increasing protein (BPI) protein products.

BPI is a protein isolated from the granules of mammalian  
polymorphonuclear leukocytes (PMNs or neutrophils), which are blood cells  
15 essential in the defense against invading microorganisms. Human BPI protein has  
been isolated from PMNs by acid extraction combined with either ion exchange  
chromatography [Elsbach, *J. Biol. Chem.*, 254:11000 (1979)] or *E. coli* affinity  
chromatography [Weiss, et al., *Blood*, 69:652 (1987)]. BPI obtained in such a  
manner is referred to herein as natural BPI and has been shown to have potent  
20 bactericidal activity against a broad spectrum of gram-negative bacteria. The  
molecular weight of human BPI is approximately 55,000 daltons (55 kD). The  
amino acid sequence of the entire human BPI protein and the nucleic acid  
sequence of DNA encoding the protein have been reported in Figure 1 of Gray et  
al., *J. Biol. Chem.*, 264:9505 (1989), incorporated herein by reference. The Gray  
25 et al. amino acid sequence is set out in SEQ ID NO: 1 hereto. U.S. Patent No.  
5,198,541 discloses recombinant genes encoding and methods for expression of  
BPI proteins, including BPI holoprotein and fragments of BPI.

BPI is a strongly cationic protein. The N-terminal half of BPI  
accounts for the high net positive charge; the C-terminal half of the molecule has a

09696709-102400  
004207-60496960

net charge of -3. [Elsbach and Weiss (1981), *supra*.] A proteolytic N-terminal fragment of BPI having a molecular weight of about 25 kD has an amphipathic character, containing alternating hydrophobic and hydrophilic regions. This N-terminal fragment of human BPI possesses the anti-bacterial efficacy of the

5 naturally-derived 55 kD human BPI holoprotein. [Ooi et al., *J. Bio. Chem.*, 262: 14891-14894 (1987)]. In contrast to the N-terminal portion, the C-terminal region of the isolated human BPI protein displays only slightly detectable anti-bacterial activity against gram-negative organisms. [Ooi et al., *J. Exp. Med.*, 174:649 (1991).]

10 "rBPI<sub>23</sub>," has been produced by recombinant means and also retains anti-bacterial activity against gram-negative organisms. Gazzano-Santoro et al., *Infect. Immun.* 60:4754-4761 (1992).

The bactericidal effect of BPI has been reported to be highly specific to gram-negative species, e.g., in Elsbach and Weiss, *Inflammation: Basic Principles and Clinical Correlates*, eds. Gallin et al., Chapter 30, Raven Press, Ltd. (1992). BPI is commonly thought to be non-toxic for other

15 microorganisms, including yeast, and for higher eukaryotic cells. Elsbach and Weiss (1992), *supra*, reported that BPI exhibits anti-bacterial activity towards a broad range of gram-negative bacteria at concentrations as low as  $10^{-8}$  to  $10^{-9}$  M, but that 100- to 1,000-fold higher concentrations of BPI were non-toxic to all of

20 the gram-positive bacterial species, yeasts, and higher eukaryotic cells tested at that time. It was also reported that BPI at a concentration of  $10^{-6}$  M or 160  $\mu$ g/ml had no toxic effect, when tested at a pH of either 7.0 or 5.5, on the gram-positive organisms *Staphylococcus aureus* (four strains), *Staphylococcus epidermidis*,

25 *Streptococcus faecalis*, *Bacillus subtilis*, *Micrococcus lysodeikticus*, and *Listeria monocytogenes*. BPI at  $10^{-6}$  M reportedly had no toxic effect on the fungi *Candida albicans* and *Candida parapsilosis* at pH 7.0 or 5.5, and was non-toxic to higher eukaryotic cells such as human, rabbit and sheep red blood cells and several human tumor cell lines. See also Elsbach and Weiss, *Advances in Inflammation Research*, ed. G. Weissmann, Vol. 2, pages 95-113 Raven Press (1981). This

30

004201" 60496960

reported target cell specificity was believed to be the result of the strong attraction of BPI for lipopolysaccharide (LPS), which is unique to the outer membrane (or envelope) of gram-negative organisms.

5 The precise mechanism by which BPI kills gram-negative bacteria is not yet completely elucidated, but it is believed that BPI must first bind to the surface of the bacteria through electrostatic and hydrophobic interactions between the cationic BPI protein and negatively charged sites on LPS. LPS has been referred to as "endotoxin" because of the potent inflammatory response that it stimulates, i.e., the release of mediators by host inflammatory cells which may  
10 ultimately result in irreversible endotoxic shock. BPI binds to lipid A, reported to be the most toxic and most biologically active component of LPS.

In susceptible gram-negative bacteria, BPI binding is thought to disrupt LPS structure, leading to activation of bacterial enzymes that degrade phospholipids and peptidoglycans, altering the permeability of the cell's outer  
15 membrane, and initiating events that ultimately lead to cell death. [Elsbach and Weiss (1992), *supra*]. BPI is thought to act in two stages. The first is a sublethal stage that is characterized by immediate growth arrest, permeabilization of the outer membrane and selective activation of bacterial enzymes that hydrolyze phospholipids and peptidoglycans. Bacteria at this stage can be rescued by growth  
20 in serum albumin supplemented media [Mannion et al., *J. Clin. Invest.*, 85:853-860 (1990)]. The second stage, defined by growth inhibition that cannot be reversed by serum albumin, occurs after prolonged exposure of the bacteria to BPI and is characterized by extensive physiologic and structural changes, including apparent damage to the inner cytoplasmic membrane.

25 Initial binding of BPI to LPS leads to organizational changes that probably result from binding to the anionic groups in the KDO region of LPS, which normally stabilize the outer membrane through binding of  $Mg^{++}$  and  $Ca^{++}$ . Attachment of BPI to the outer membrane of gram-negative bacteria produces rapid permeabilization of the outer membrane to hydrophobic agents such as  
30 actinomycin D. Binding of BPI and subsequent gram-negative bacterial killing

09696709-102400

depends, at least in part, upon the LPS polysaccharide chain length, with long O-chain bearing, "smooth" organisms being more resistant to BPI bactericidal effects than short O-chain bearing, "rough" organisms [Weiss et al., *J. Clin. Invest.* 65: 619-628 (1980)]. This first stage of BPI action, permeabilization of the gram-negative outer envelope, is reversible upon dissociation of the BPI, a process requiring the presence of divalent cations and synthesis of new LPS [Weiss et al., *J. Immunol.* 132: 3109-3115 (1984)]. Loss of gram-negative bacterial viability, however, is not reversed by processes which restore the envelope integrity, suggesting that the bactericidal action is mediated by additional lesions induced in the target organism and which may be situated at the cytoplasmic membrane (Mannion et al., *J. Clin. Invest.* 86: 631-641 (1990)). Specific investigation of this possibility has shown that on a molar basis BPI is at least as inhibitory of cytoplasmic membrane vesicle function as polymyxin B (In't Veld et al., *Infection and Immunity* 56: 1203-1208 (1988)) but the exact mechanism as well as the relevance of such vesicles to studies of intact organisms has not yet been elucidated.

BPI is also capable of neutralizing the endotoxic properties of LPS to which it binds. Because of its bactericidal properties for gram-negative organisms and its ability to neutralize LPS, BPI can be utilized for the treatment of mammals suffering from diseases caused by gram-negative bacteria, such as bacteremia or sepsis.

Poloxamer (polyoxypropylene-polyoxyethylene block copolymer) surfactants are non-ionic block copolymer surfactants having a structure composed of two blocks or chains of hydrophilic polyoxyethylene (POE) flanking a single block of hydrophobic polyoxypropylene (POP). They are considered to be among the least toxic of known surfactants and are widely used in foods, drugs and cosmetics.

Of interest to the present invention is co-owned, co-pending allowed U.S. Patent Application Serial No. 08/190,869 (PCT Application Publication No. WO 94/17819), herein incorporated by reference, which describes the improved

00696709-102400

solubilization or stability of pharmaceutical compositions containing BPI protein products and a poloxamer surfactant, either alone or in combination with a polysorbate surfactant.

Also of interest to the present invention are PCT Application  
5 Publication No. WO88/06038 and U.S. Patent No. 5,183,687, which address use of poloxamer surfactants with and without "conventional" antibiotics in the treatment of viral, *Mycobacterium* and *Coccidioides* infections.

There exists a desire in the art for methods and compositions  
capable of improving the therapeutic effectiveness of antibacterial agents such as  
10 BPI protein products. Such methods and compositions could ideally reduce the dosage of agent required to achieve desired therapeutic effects.

#### SUMMARY OF THE INVENTION

The present invention provides improved anti-microbial compositions and methods of treatment. According to one aspect of the invention,  
15 improved therapeutic compositions are provided that comprise a BPI protein product and a polyoxypropylene-polyoxyethylene block copolymer (poloxamer) surfactant that enhances the anti-bacterial activity of the BPI protein product. Presently preferred bactericidal-activity-enhancing poloxamer surfactants include poloxamer 333 (PLURONIC 103, BASF, Parsippany, NJ), poloxamer 334  
20 (PLURONIC 104, BASF), poloxamer 335 (PLURONIC 105, BASF), or poloxamer 403 (PLURONIC P123, BASF). Poloxamers employed according to the invention may optionally be heat-treated prior to incorporation into the compositions. Especially preferred are compositions including poloxamer 333 or poloxamer 403. This aspect of the invention is based upon the finding that the  
25 combination of a BPI protein product with one of the above-listed poloxamer surfactants unexpectedly enhances the bactericidal activity of the BPI protein product, both *in vitro* and *in vivo*. The improved therapeutic compositions of the present invention may further comprise ethylenediaminetetraacetic acid (EDTA). This aspect of the invention is based on the discovery that the addition of EDTA

004201 60296960  
09696709 102400

to therapeutic compositions containing BPI protein product and a bactericidal-activity-enhancing poloxamer surfactant (such as poloxamer 333, poloxamer 334, poloxamer 335 or poloxamer 403) may produce further enhancement of the bactericidal activity of the BPI protein product.

5 Corresponding improved methods for treating bacterial infection are also provided; the improvement comprising administering to a patient with a suspected or confirmed infection a therapeutic composition of BPI protein product and a bactericidal-activity-enhancing poloxamer, and optionally EDTA. The present invention also contemplates the use of a bactericidal-activity-enhancing  
10 poloxamer surfactant (such as poloxamer 333, poloxamer 334, poloxamer 335, or poloxamer 403) with a BPI protein product, and optionally EDTA, for the manufacture of a medicament for treatment of bacterial infection.

The present invention further provides improved compositions for inhibiting bacterial and fungal growth comprising a BPI protein product and a  
15 bacterial and fungal growth-inhibiting enhancing poloxamer surfactant, and optionally EDTA. This aspect of the invention is based upon the discovery that combination of a BPI protein product with a bacterial and fungal growth-inhibiting enhancing poloxamer surfactant unexpectedly enhances the growth-inhibitory activity of the BPI protein product. Corresponding methods of killing or inhibiting  
20 the growth of bacteria or fungi are provided that comprise contacting the organisms with a composition comprising a BPI protein product and a bacterial and fungal growth-inhibiting enhancing poloxamer surfactant, and optionally EDTA. Presently preferred bacterial and fungal growth-inhibiting enhancing poloxamer surfactants include poloxamer 333, poloxamer 334, poloxamer 335, and  
25 poloxamer 403.

With regard to the improved methods for treating bacterial infection described above, a method of improving the therapeutic effectiveness of antibiotics for treatment of bacterial infections is also provided. According to this method, the antibiotic is concurrently administered with a composition comprising a BPI  
30 protein product formulated with a BPI-activity-enhancing poloxamer surfactant

004201 60296960

00420T" 60296969

(such as poloxamer 333, poloxamer 334, poloxamer 335, or poloxamer 403), and optionally with EDTA. This aspect of the invention is based on the discovery that the improvement in therapeutic effectiveness of antibiotics that is seen with the addition of BPI protein product can be further enhanced by various poloxamer formulations, and that the addition of EDTA to the BPI protein product/poloxamer formulation provides an even greater enhancement of the antibiotic's therapeutic effectiveness. This aspect of the invention also provides use of poloxamer surfactants (such as poloxamer 333, poloxamer 334, poloxamer 335, or poloxamer 403), optionally with EDTA, for the manufacture of a medicament containing BPI protein product for co-treatment of a bacterial infection with an antibiotic.

The following findings are illustrative of this aspect of the invention: For a *Pseudomonas* species, enhancement of the improved therapeutic effectiveness of ceftizoxime was provided by BPI protein product formulations containing poloxamer 333, poloxamer 335, or poloxamer 403; enhancement for ceftriaxone was provided by BPI protein product formulations containing poloxamer 333, poloxamer 335, or poloxamer 403; and enhancement for chloramphenicol was provided by BPI protein product formulations containing poloxamer 333, poloxamer 334, poloxamer 335, or poloxamer 403. For an *Acinetobacter* species, enhancement for ceftazidime was provided by BPI protein product formulations containing poloxamer 333, poloxamer 334, poloxamer 335, or poloxamer 403; enhancement for ceftriaxone was provided by BPI protein product formulations containing poloxamer 333, poloxamer 334, poloxamer 335, or poloxamer 403; and enhancement for chloramphenicol was provided by BPI protein product formulations containing poloxamer 333, poloxamer 334, poloxamer 335, or poloxamer 403. For a *Streptococcus* species, enhancement for oxacillin was provided by BPI protein product formulations containing poloxamer 333, poloxamer 334, poloxamer 335, or poloxamer 403. For an *Enterococcus* species, enhancement for rifampicin was provided by BPI protein product formulations containing poloxamer 335 or poloxamer 403; and enhancement for



ciprofloxacin was provided by BPI protein product formulations containing poloxamer 333.

For a *Pseudomonas* species, enhancement of the therapeutic effectiveness of a variety of antibiotics was provided by a BPI protein product formulation containing poloxamer 403, and even greater enhancement was provided by adding increasing concentrations of EDTA to the BPI/poloxamer 403 formulation.

Numerous additional aspects and advantages of the invention will become apparent to those skilled in the art upon consideration of the following detailed description of the invention which describes presently preferred embodiments thereof.

#### DETAILED DESCRIPTION

The present invention provides improved anti-microbial compositions and methods of treatment. The improved methods and compositions, in addition to being useful for treatment of bacterial infections and conditions associated therewith or resulting therefrom (such as sepsis and bacteremia), and are also useful for prophylaxis of patients at high risk of bacterial infection, e.g., patients who will undergo abdominal or genitourinary surgery, or trauma victims.

Specifically, the present invention provides, in a therapeutic composition comprising a BPI protein product and a stabilizing poloxamer surfactant, the improvement comprising a bactericidal-activity-enhancing poloxamer surfactant, such as poloxamer 333, poloxamer 334, poloxamer 335, or poloxamer 403. The present invention is based upon the finding that the combination of a BPI protein product with one of these above-listed poloxamer surfactants unexpectedly enhances the bactericidal activity of the BPI protein product, both *in vitro* and *in vivo*. The improved therapeutic compositions of the present invention may further comprise EDTA. This aspect of the invention is based on the discovery that the addition of EDTA to some therapeutic compositions containing BPI protein product and a bactericidal-activity-enhancing

004201" 60296960

poloxamer surfactant, such as poloxamer 333, poloxamer 334, poloxamer 335, or poloxamer 403, produces further enhancement of the bactericidal activity of the BPI protein product. Such compositions may optionally comprise pharmaceutically acceptable diluents, adjuvants or carriers. The invention utilizes  
5 any of the large variety of BPI protein products known to the art including natural BPI protein, recombinant BPI protein, BPI fragments, BPI analogs, BPI variants, and BPI peptides.

Corresponding improved methods for treating bacterial infection are also provided, the improvement comprising administering to a patient with a  
10 suspected or confirmed infection a therapeutic composition of BPI protein product and a bactericidal-activity-enhancing poloxamer, and optionally EDTA. The present invention also contemplates the use of a bactericidal-activity-enhancing poloxamer surfactant (such as poloxamer 333, poloxamer 334, poloxamer 335, or poloxamer 403) with a BPI protein product, and optionally EDTA, for the  
15 manufacture of a medicament for treatment of bacterial infection. The therapeutic composition of BPI protein product and poloxamer surfactant with or without EDTA may be administered systemically or topically to a subject suffering from a suspected or confirmed bacterial infection.

Poloxamer 333 is sold by BASF (Parsippany, NJ) under the name  
20 PLURONIC P103 and has a molecular weight of 4950 and a hydrophilic/lipophilic balance (HLB) value of 7-12. Poloxamer 334 is sold by BASF under the name PLURONIC P104 and has a molecular weight of 5900 and an HLB value of 12-18. Poloxamer 335 is sold by BASF under the name PLURONIC P105 and has a molecular weight of 6500 and an HLB value of 12-18. Poloxamer 403 is sold by  
25 BASF under the name PLURONIC P123 and has a molecular weight of 5750 and an HLB value of 7-12. Presently preferred bactericidal-activity-enhancing poloxamer surfactants include poloxamer 333, poloxamer 334, poloxamer 335 or poloxamer 403. Especially preferred are compositions including poloxamer 333 or poloxamer 403.

004201" 60296960  
09696709 102400

Poloxamers employed according to the invention may optionally be heat-treated prior to incorporation into the compositions. A preferred method of heat treatment is as follows: (1) making a solution of the poloxamer in deionized water, (2) heating the solution to a boil, (3) removing it from heat, (4) allowing it to cool to room temperature, and (5) stirring until the poloxamer is completely solubilized. Alternatively, in the heating step (2), the solution may be boiled for up to 30 minutes or more.

The present invention further provides improved compositions for inhibiting bacterial and fungal growth comprising a BPI protein product and a bacterial and fungal growth-inhibiting enhancing poloxamer surfactant, and optionally EDTA. This aspect of the invention is based upon the discovery that a bacterial and fungal growth-inhibiting enhancing poloxamer surfactant unexpectedly enhances the growth-inhibitory activity of BPI protein product, and that improved compositions comprising such poloxamer surfactants and BPI protein product display superior growth-inhibitory preservative effects. Corresponding methods of killing or inhibiting the growth of bacteria or fungi are provided that comprise contacting the organisms with a composition comprising a BPI protein product and a bacterial and fungal growth-inhibiting enhancing poloxamer surfactant, and optionally EDTA. Presently preferred bacterial and fungal growth-inhibiting enhancing poloxamer surfactants include poloxamer 333, poloxamer 334, poloxamer 335, and poloxamer 403.

These methods can be practiced *in vivo* or in a variety of *in vitro* uses such as use as a preservative, use to decontaminate fluids and surfaces, or use to sterilize surgical and other medical equipment and implantable devices, including prosthetic joints. These methods can also be used for *in situ* sterilization of indwelling invasive devices such as intravenous lines and catheters which are often foci of infection and in the preparation of growth media for cells. The efficacy of the improved compositions for inhibiting bacterial and fungal growth can be evaluated according to the assay described below in Example 8, or by any of the assays described in co-owned, copending patent application Cohen et al.,

004207" 60296960

U.S. Serial No. 08/125,651 filed September 22, 1993, and continuation-in-part thereof U.S. Serial No. 08/273,401 filed July 11, 1994, and continuation-in-part thereof U.S. Serial No. 08/311,611 filed September 22, 1994, and corresponding PCT Application No. PCT/US94/11225, and co-owned, copending patent application (Little et al.) U.S. Serial No. 08/183,222 filed January 14, 1994, and continuation-in-part thereof U.S. Serial No. 08/209,762 filed March 11, 1994, and continuation-in-part thereof (Horwitz et al.) U.S. Serial No. 08/274,299 filed July 11, 1994, and continuation-in-part thereof U.S. Serial No. 08/372,783 filed January 13, 1995, and corresponding PCT Application No. PCT/US95/00656, and co-owned, copending patent application Little et al., U.S. Serial No. 08/183,222 filed January 14, 1994, and continuation-in-part thereof U.S. Serial No. 08/209,762 filed March 11, 1994, and continuation-in-part thereof U.S. Serial No. 08/273,540 filed July 11, 1994, and continuation-in-part thereof U.S. Serial No. 08/372,105 filed January 13, 1995, and corresponding PCT Application No. PCT/US95/00498, all of which are incorporated herein by reference.

BPI protein product is thought to interact with a variety of host defense elements present in whole blood or serum, including complement, p15 and LBP, and other cells and components of the immune system. Such interactions may result in potentiation of the activities of BPI protein product. Because of these interactions, BPI protein products can be expected to exert even greater activity *in vivo* than *in vitro*. Thus, while *in vitro* tests are predictive of *in vivo* utility, absence of activity *in vitro* does not necessarily indicate absence of activity *in vivo*. For example, BPI has been observed to display a greater bactericidal effect on gram-negative bacteria in whole blood or plasma assays than in assays using conventional media. [Weiss et al., *J. Clin. Invest.* 90:1122-1130 (1992)]. This is also shown in *in vivo* animal experiments (see, e.g., co-owned, copending U.S. Application Cohen et al., U.S. Serial NO. 08/311,611 filed September 22, 1994, and corresponding PCT Appl. No. PCT/US94/11225, all of which are incorporated herein by reference. This may be because conventional *in vitro* systems lack the blood elements that facilitate or potentiate BPI's function *in vivo*,

004201" 60296960

or because conventional media contain higher than physiological concentrations of magnesium and calcium, which are typically inhibitors of the anti-bacterial activity of BPI protein products. Furthermore, in the host, BPI protein product is available to neutralize endotoxin released during host infection, including from stress-induced translocation of gram-negative bacteria or from antibiotic treatment of gram-negative bacteria, a further clinical benefit not seen in or predicted by *in vitro* tests.

It is also contemplated that the BPI protein product be administered with other products that potentiate the bactericidal activity of BPI protein products. For example, serum complement potentiates the gram-negative bactericidal activity of BPI protein products; the combination of BPI protein product and serum complement provides synergistic bactericidal/growth inhibitory effects. See, e.g., Ooi *et al. J. Biol. Chem.*, 265: 15956 (1990) and Levy *et al. J. Biol. Chem.*, 268: 6038-6083 (1993) which address naturally-occurring 15 kD proteins potentiating BPI antibacterial activity. See also co-owned, co-pending PCT Application No. US94/07834 filed July 13, 1994, which corresponds to U.S. Patent Application Serial No. 08/274,303 filed July 11, 1994 as a continuation-in-part of U.S. Patent Application Serial No. 08/093,201 filed July 14, 1993. These applications, which are all incorporated herein by reference, describe methods for potentiating gram-negative bactericidal activity of BPI protein products by administering lipopolysaccharide binding protein (LBP) and LBP protein products. LBP protein derivatives and derivative hybrids which lack CD-14 immunostimulatory properties are described in PCT Application No. US94/06931 filed June 17, 1994, which corresponds to co-owned, co-pending U.S. Patent Application Serial No. 08/261,660, filed June 17, 1994 as a continuation-in-part of U.S. Patent Application Serial No. 08/079,510, filed June 17, 1993, the disclosures of all of which are hereby incorporated by reference.

An advantage provided by the present invention is the ability to provide more effective killing or growth inhibition of bacteria and fungi and enhanced anti-bacterial or anti-fungal activity of the BPI protein product.

004207" 60796960

Therapeutic compositions comprising BPI protein product and a BPI anti-microbial activity enhancing poloxamer surfactant, and optionally containing EDTA, may be administered systemically or topically. Systemic routes of administration include oral, intravenous, intramuscular or subcutaneous injection (including into a depot for long-term release), intraocular and retrobulbar, intrathecal, intraperitoneal (e.g. by intraperitoneal lavage), transpulmonary using aerosolized or nebulized drug, or transdermal. For example, when given parenterally, BPI protein product compositions are generally injected in doses ranging from 1  $\mu\text{g/kg}$  to 100  $\text{mg/kg}$  per day, and preferably at doses ranging from 0.1  $\text{mg/kg}$  to 20  $\text{mg/kg}$  per day. The treatment may continue at the same, reduced or increased dose per day for, e.g., 1 to 3 days, and additionally as determined by the treating physician. Topical routes include administration in the form of salves, ophthalmic drops, ear drops, irrigation fluids (for, e.g., irrigation of wounds) or medicated shampoos. For example, for topical administration in drop form, about 10 to 200  $\mu\text{L}$  of a BPI protein product composition may be applied one or more times per day as determined by the treating physician. Those skilled in the art can readily optimize effective dosages and administration regimens for therapeutic compositions comprising BPI protein product and a BPI bactericidal-activity enhancing poloxamer surfactant, and optionally containing EDTA, as determined by good medical practice and the clinical condition of the individual patient.

With regard to the improved methods for treating bacterial infection described above, a method of improving the therapeutic effectiveness of antibiotics for treatment of bacterial infections is also provided. According to this method, the antibiotic is concurrently administered with a composition comprising a BPI protein product formulated with a BPI-activity-enhancing poloxamer surfactant (such as poloxamer 333, poloxamer 334, poloxamer 335, or poloxamer 403), and optionally with EDTA. This aspect of the invention is based on the discovery that the improvement in therapeutic effectiveness of antibiotics that is seen with the addition of BPI protein product can be further enhanced by various poloxamer formulations, and that the addition of EDTA to the BPI protein product/poloxamer

formulation provides an even greater enhancement of the antibiotic's therapeutic effectiveness. This aspect of the invention also provides use of poloxamer surfactants (such as poloxamer 333, poloxamer 334, poloxamer 335, or poloxamer 403), optionally with EDTA, for the manufacture of a medicament containing BPI protein product for co-treatment of a bacterial infection with an antibiotic.

For this aspect of the invention, the improved therapeutic effectiveness of antibiotics seen upon concurrent administration with BPI protein product can be observed in a number of ways. For example, a BPI protein product may convert an organism that is clinically resistant to an antibiotic into an organism that is clinically susceptible to the antibiotic, or may otherwise improve the antibiotic susceptibility of that organism. The BPI protein product and antibiotic may have a therapeutic effect when both are given in doses below the amounts sufficient for monotherapeutic effectiveness. The inclusion of a BPI-activity-enhancing poloxamer surfactant in the BPI protein product formulation provides a further enhancement of these activities. Co-owned, copending patent application Cohen et al., U.S. Serial No. 08/125,651 filed September 22, 1993, and continuation-in-part thereof U.S. Serial No. 08/273,401 filed July 11, 1994, and continuation-in-part thereof U.S. Serial No. 08/311,611 filed September 22, 1994, and corresponding PCT Application No. PCT/US94/11225, and co-owned, copending patent application (Little et al.), U.S. Serial No. 08/183,222 filed January 14, 1994, and continuation-in-part thereof U.S. Serial No. 08/209,762 filed March 11, 1994, and continuation-in-part thereof (Horwitz et al.) U.S. Serial No. 08/274,299 filed July 11, 1994, and continuation-in-part thereof U.S. Serial No. 08/372,783 filed January 13, 1995, and corresponding PCT Application No. PCT/US95/00656, all of which are incorporated herein by reference, disclose methods for evaluating the use of BPI as an anti-microbial agent and to enhance the effectiveness of antibiotics.

The improved therapeutic effectiveness of antibiotics may be demonstrated in *in vivo* animal models, or may be predicted on the basis of a variety of *in vitro* tests, including (1) determinations of the minimum inhibitory

004207" 60296660

concentration (MIC) of an antibiotic required to inhibit growth of a gram-negative organism for 24 hours, (2) determinations of the effect of an antibiotic on the kinetic growth curve of a gram-negative organism, and (3) checkerboard assays of the MIC of serial dilutions of antibiotic alone or in combination with serial dilutions of BPI protein product. Such improved effectiveness may be demonstrated by (a) a reduction in the number of organisms, (b) a reduced MIC, and/or (c) reversal of the organism's resistance to the antibiotic. Exemplary models or tests are described in Eliopoulos and Moellering In *Antibiotics in Laboratory Medicine*, 3rd ed. (Lorian, V., Ed.) pp. 432-492, Williams and Wilkins, Baltimore MD (1991).

"Concurrent administration," or co-treatment, as used herein includes administration of the agents, in conjunction or combination, together, or before or after each other. The BPI protein product (formulated with activity-enhancing poloxamer) and antibiotics may be administered by different routes. For example, the formulated BPI protein product may be administered intravenously while the antibiotics are administered intramuscularly, intravenously, subcutaneously, orally or intraperitoneally. Alternatively, the formulated BPI protein product may be administered intraperitoneally while the antibiotics are administered intraperitoneally or intravenously, or the formulated BPI protein product may be administered in an aerosolized or nebulized form while the antibiotics are administered, e.g., intravenously. The formulated BPI protein product and antibiotics are preferably both administered intravenously. The formulated BPI protein product and antibiotics may be given sequentially in the same intravenous line, after an intermediate flush, or may be given in different intravenous lines. The formulated BPI protein product and antibiotics may be administered simultaneously or sequentially, as long as they are given in a manner sufficient to allow both agents to achieve effective concentrations at the site of infection.

Concurrent administration of formulated BPI protein product and antibiotic is expected to provide more effective treatment of bacterial infections.

09596709-102400



Concurrent administration of the two agents may provide greater therapeutic effects *in vivo* than either agent provides when administered singly. It may permit a reduction in the dosage of one or both agents with achievement of a similar therapeutic effect. Alternatively, the concurrent administration may produce a more rapid or complete bactericidal/bacteriostatic effect than could be achieved with either agent alone.

Therapeutic effectiveness is correlated with a successful clinical outcome, and does not require that the antimicrobial agent or agents kill 100% of the organisms involved in the infection. Success depends on achieving a level of antibacterial activity at the site of infection that is sufficient to inhibit the bacteria in a manner that tips the balance in favor of the host. When host defenses are maximally effective, the antibacterial effect required may be minimal. Reducing organism load by even one log (a factor of 10) may permit the host's own defenses to control the infection. In addition, augmenting an early bactericidal/bacteriostatic effect can be more important than long-term bactericidal/bacteriostatic effect. These early events are a significant and critical part of therapeutic success, because they allow time for host defense mechanisms to activate. Increasing the bactericidal rate may be particularly important for infections such as meningitis, bone or joint infections [Stratton, *Antibiotics in Laboratory Medicine*, 3rd ed. (Lorian, V., Ed.) pp. 849-879, Williams and Wilkins, Baltimore MD (1991)], or alternatively, for infections involving slow-growing organisms which may have a decreased sensitivity to antibiotics.

As used herein, "BPI protein product" includes naturally and recombinantly produced BPI protein; natural, synthetic, and recombinant biologically active polypeptide fragments of BPI protein; biologically active polypeptide variants of BPI protein or fragments thereof, including hybrid fusion proteins and dimers; biologically active polypeptide analogs of BPI protein or fragments or variants thereof, including cysteine-substituted analogs; and BPI-derived peptides. The BPI protein products administered according to this invention may be generated and/or isolated by any means known in the art. U.S.

09696709 "102400

Patent No. 5,198,541, the disclosure of which is incorporated herein by reference, discloses recombinant genes encoding and methods for expression of BPI proteins including recombinant BPI holoprotein, referred to as rBPI<sub>50</sub> and recombinant fragments of BPI. Co-owned, copending U.S. Patent Application Ser. No.

- 5 07/885,501 and a continuation-in-part thereof, U.S. Patent Application Ser. No. 08/072,063 filed May 19, 1993 and corresponding PCT Application No. 93/04752 filed May 19, 1993, which are all incorporated herein by reference, disclose novel methods for the purification of recombinant BPI protein products expressed in and secreted from genetically transformed mammalian host cells in culture and
- 10 discloses how one may produce large quantities of recombinant BPI products suitable for incorporation into stable, homogeneous pharmaceutical preparations.

- Biologically active fragments of BPI (BPI fragments) include biologically active molecules that have the same or similar amino acid sequence as a natural human BPI holoprotein, except that the fragment molecule lacks amino-
- 15 terminal amino acids, internal amino acids, and/or carboxy-terminal amino acids of the holoprotein. Nonlimiting examples of such fragments include a N-terminal fragment of natural human BPI of approximately 25 kD, described in Ooi et al., *J. Exp. Med.*, 174:649 (1991), and the recombinant expression product of DNA encoding N-terminal amino acids from 1 to about 193 or 199 of natural human
- 20 BPI, described in Gazzano-Santoro et al., *Infect. Immun.* 60:4754-4761 (1992), and referred to as rBPI<sub>23</sub>. In that publication, an expression vector was used as a source of DNA encoding a recombinant expression product (rBPI<sub>23</sub>) having the 31-residue signal sequence and the first 199 amino acids of the N-terminus of the mature human BPI, as set out in Figure 1 of Gray et al., *supra*, except that valine
- 25 at position 151 is specified by GTG rather than GTC and residue 185 is glutamic acid (specified by GAG) rather than lysine (specified by AAG). Recombinant holoprotein (rBPI) has also been produced having the sequence (SEQ ID NOS: 1 and 2) set out in Figure 1 of Gray et al., *supra*, with the exceptions noted for rBPI<sub>23</sub> and with the exception that residue 417 is alanine (specified by GCT) rather
- 30 than valine (specified by GTT). Other examples include dimeric forms of BPI

004207 60296960

fragments, as described in co-owned and co-pending U.S. Patent Application Serial No. 08/212,132, filed March 11, 1994, and corresponding PCT Application No. PCT/US95/03125, the disclosures of which are incorporated herein by reference. Preferred dimeric products include dimeric BPI protein products wherein the monomers are amino-terminal BPI fragments having the N-terminal residues from about 1 to 175 to about 1 to 199 of BPI holoprotein. A particularly preferred dimeric product is the dimeric form of the BPI fragment having N-terminal residues 1 through 193, designated rBPI<sub>42</sub> dimer.

Biologically active variants of BPI (BPI variants) include but are not limited to recombinant hybrid fusion proteins, comprising BPI holoprotein or biologically active fragment thereof and at least a portion of at least one other polypeptide, and dimeric forms of BPI variants. Examples of such hybrid fusion proteins and dimeric forms are described by Theofan et al. in co-owned, copending U.S. Patent Application Serial No. 07/885,911, and a continuation-in-part application thereof, U.S. Patent Application Serial No. 08/064,693 filed May 19, 1993 and corresponding PCT Application No. US93/04754 filed May 19, 1993, which are all incorporated herein by reference and include hybrid fusion proteins comprising, at the amino-terminal end, a BPI protein or a biologically active fragment thereof and, at the carboxy-terminal end, at least one constant domain of an immunoglobulin heavy chain or allelic variant thereof. Similarly configured hybrid fusion proteins involving part or all Lipopolysaccharide Binding Protein (LBP) are also contemplated for use in the present invention.

Biologically active analogs of BPI (BPI analogs) include but are not limited to BPI protein products wherein one or more amino acid residues have been replaced by a different amino acid. For example, co-owned, copending U.S. Patent Application Ser. No. 08/013,801 filed February 2, 1993 and corresponding PCT Application No. US94/01235 filed February 2, 1994, the disclosures of which are incorporated herein by reference, discloses polypeptide analogs of BPI and BPI fragments wherein a cysteine residue is replaced by a different amino acid. A preferred BPI protein product described by this application is the

004201 60296960

expression product of DNA encoding from amino acid 1 to approximately 193 or 199 of the N-terminal amino acids of BPI holoprotein, but wherein the cysteine at residue number 132 is substituted with alanine and is designated rBPI<sub>21</sub>Δcys or rBPI<sub>21</sub>. Other examples include dimeric forms of BPI analogs; e.g. co-owned and co-pending U.S. Patent Application Serial No. 08/212,132 filed March 11, 1994, and corresponding PCT Application No. PCT/US95/03125, the disclosures of which are incorporated herein by reference.

Other BPI protein products useful according to the methods of the invention are peptides derived from or based on BPI produced by recombinant or synthetic means (BPI-derived peptides), such as those described in co-owned and co-pending U.S. Patent Application Serial No. 08/504,841 filed July 20, 1995 and in co-owned and copending PCT Application No. PCT/US94/10427 filed September 15, 1994, which corresponds to U.S. Patent Application Serial No. 08/306,473 filed September 15, 1994, and PCT Application No. US94/02465 filed March 11, 1994, which corresponds to U.S. Patent Application Serial No. 08/209,762, filed March 11, 1994, which is a continuation-in-part of U.S. Patent Application Serial No. 08/183,222, filed January 14, 1994, which is a continuation-in-part of U.S. Patent Application Ser. No. 08/093,202 filed July 15, 1993 (for which the corresponding international application is PCT Application No. US94/02401 filed March 11, 1994), which is a continuation-in-part of U.S. Patent Application Ser. No. 08/030,644 filed March 12, 1993, the disclosures of all of which are incorporated herein by reference.

Presently preferred BPI protein products include recombinantly-produced N-terminal fragments of BPI, especially those having a molecular weight of approximately between 21 to 25 kD such as rBPI<sub>23</sub> or rBPI<sub>21</sub>, or dimeric forms of these N-terminal fragments (e.g., rBPI<sub>42</sub> dimer). Additionally, preferred BPI protein products include rBPI<sub>50</sub> and BPI-derived peptides.

The administration of BPI protein products is preferably accomplished with a pharmaceutical composition comprising a BPI protein product and a pharmaceutically acceptable diluent, adjuvant, or carrier. The BPI protein

004201" 6029696

product may be administered without or in conjunction with known surfactants, other chemotherapeutic agents or additional known anti-microbial agents. One pharmaceutical composition containing BPI protein products (e.g., rBPI<sub>50</sub>, rBPI<sub>23</sub>) comprises the BPI protein product at a concentration of 1 mg/ml in citrate buffered saline (5 or 20 mM citrate, 150 mM NaCl, pH 5.0) comprising 0.1% by weight of poloxamer 188 (Pluronic F-68, BASF Wyandotte, Parsippany, NJ) and 0.002% by weight of polysorbate 80 (Tween 80, ICI Americas Inc., Wilmington, DE). Another pharmaceutical composition containing BPI protein products (e.g., rBPI<sub>21</sub>) comprises the BPI protein product at a concentration of 2 mg/mL in 5 mM citrate, 150 mM NaCl, 0.2% poloxamer 188 and 0.002% polysorbate 80. Such combinations are described in co-owned, co-pending PCT Application No. US94/01239 filed February 2, 1994, which corresponds to U.S. Patent Application Ser. No. 08/190,869 filed February 2, 1994 and U.S. Patent Application Ser. No. 08/012,360 filed February 2, 1993, the disclosures of all of which are incorporated herein by reference.

Other aspects and advantages of the present invention will be understood upon consideration of the following illustrative examples. Example 1 addresses the effects of poloxamer 403 or poloxamer 334 on the bactericidal activity of BPI protein products against *S. aureus* or *A. baumannii* (formerly *A. anitratus*) in water. Example 2 addresses the effects of poloxamer 333 or poloxamer 403 on the bactericidal activity of non-formulated or formulated BPI protein products against *A. baumannii*, *S. aureus*, *N. meningitidis* or *P. aeruginosa* in serum, broth or water. Example 3 addresses the effects of poloxamer 333 or poloxamer 334 on the bactericidal activity of BPI protein products against *S. pneumoniae*, *S. aureus*, *E. faecium*, or *A. baumannii* in water. Example 4 relates to uses of other poloxamers. Example 5 addresses the effects of poloxamers 188, 333, 334, 335, or 403 (with or without EDTA) on the bactericidal activity of BPI protein products against *A. baumannii*, *S. aureus*, *S. pneumoniae*, *E. faecium*, or *P. aeruginosa* in serum, Mueller-Hinton broth, tryptic soy broth, or water. Example 6 addresses the effect of compositions containing

004207 60796960  
09696709 102400

BPI protein product and poloxamer 188, 333, 334, 335, or 403 in the presence or absence of EDTA on the susceptibility of a variety of organisms to antibiotics. Example 7 addresses the effect of compositions containing BPI protein product and an anti-bacterial activity-enhancing poloxamer surfactant in a rabbit model of corneal injury and ulceration. Example 8 addresses the effect of compositions containing BPI protein product and poloxamer 188 or 403 in the presence or absence of EDTA on the growth of various bacteria and fungi.

#### EXAMPLE 1

##### 10 BACTERICIDAL ACTIVITY OF COMPOSITIONS CONTAINING BPI PROTEIN PRODUCT AND POLOXAMER 403 OR POLOXAMER 334 ON *S. AUREUS* AND *A. BAUMANNII* IN WATER

The bactericidal activity of therapeutic compositions comprising BPI protein product and either poloxamer 403 (PLURONIC P123, BASF Wyandotte Corp., Parsippany, NJ), heat-treated PLURONIC 123, or heat-treated poloxamer 334 (PLURONIC P104, BASF Wyandotte Corp.), was evaluated against clinical isolates of bacteria from the Microscan® library (Dade Microscan, West Sacramento, CA). Therapeutic compositions comprising 1 mg/mL rBPI<sub>21</sub> and 0.1% (w/v) PLURONIC P123, or heat-treated PLURONIC P123, were formulated by diluting a 2 mg/mL solution of "non-formulated" rBPI<sub>21</sub> (in buffer comprising 5 mM sodium citrate and 150 mM NaCl, without any surfactants) at a 1:2 ratio with a 0.2% solution of the PLURONIC P123. A therapeutic composition comprising 2 mg/mL rBPI<sub>21</sub> and 0.1% (w/v) heat-treated PLURONIC P104 was prepared. Poloxamer control solutions containing only 0.1% PLURONIC P123 or 0.1% heat-treated PLURONIC P123, and no rBPI<sub>21</sub>, were also prepared.

25 Sterile stock solutions of 1.0% PLURONIC P123 were prepared by stirring the PLURONIC P123 in deionized water until dissolved and filtering the solution through a 0.22µm Nalgene filter unit (Nalge Co., Rochester, NY). Sterile stock solutions of heat-treated PLURONIC P123 were prepared using the following procedure: (1) making a 1.0% (w/v) solution of PLURONIC P123 in deionized water, (2) heating the solution to a boil, (3) removing it from heat, (4)

30

004207" 60796960

allowing it to cool to room temperature, (5) stirring until the PLURONIC P123 was completely solubilized, and (6) filtering the solution through a 0.22 $\mu$ m Nalgene filter unit for sterilization. Alternatively, the stock solutions may be autoclaved for sterilization. Heat-treated PLURONIC P104 was prepared similarly.

The bacteria to be used in the assays, *S. aureus* (Microscan® ID no. 052-106) and *A. baumannii* (Microscan® ID no. 12291), were grown on tryptic soy agar (TSA) plates (Remel, Catalog #01-920, Lenexa, KN) for 24 hours. A bacterial stock emulsion of about 4 to 7 x 10<sup>4</sup> cells/mL was prepared by emulsifying bacterial colonies in sterile water for injection (Kendall McGaw Laboratory, Irvine, CA) to a 0.5 McFarland standard and diluting further by 1:10 in water. Assays were conducted by adding 944  $\mu$ L of sterile water for injection to 4.5 mL polypropylene tubes (Nalgene Cryovial, Nalge Co., Rochester, NY), followed by 40  $\mu$ L of the bacterial emulsion, followed by 16  $\mu$ L of the 1 mg/mL rBPI<sub>21</sub>/0.1% PLURONIC P123 therapeutic composition or poloxamer control solution (or 8  $\mu$ L of the 2 mg/mL rBPI<sub>21</sub>/0.1% PLURONIC P104 therapeutic composition). The tubes were mixed by inversion and incubated at 37°C for 30 minutes. Following incubation, the remaining colony forming units (CFU) were counted at a 10<sup>-2</sup> dilution by plating 10 $\mu$ L from each tube onto TSA plates, and at 10<sup>-4</sup> dilutions by plating a 1:100 dilution of 10 $\mu$ L from each tube onto TSA plates. The TSA plates were incubated at 37°C for 18 hours and the number of bacterial colonies were visually counted. Results are shown below in Tables 1 and 2.

09696709-102400  
004207-60496960

Table 1

|   | <i>S. aureus</i>   | CFU    |
|---|--|--------|
|   | Positive Control   | 150000 |
|   | 16 µg/mL rBPI <sub>21</sub> with 0.1% PLURONIC P123              | 26600  |
| 5 | 16 µg/mL rBPI <sub>21</sub> with 0.1% heat-treated PLURONIC P123 | 26400  |
|   | 0.1% PLURONIC P123 control                                       | 150000 |
|   | 0.1% heat-treated PLURONIC P123 control                          | 150000 |
|   | 16 µg/mL rBPI <sub>21</sub> with 0.1% heat-treated PLURONIC P104 | 49100  |

Table 2

|    |  |       |
|----|--|-------|
|    | <i>A. baumannii</i>  | CFU   |
| 10 | Positive Growth Control (no rBPI <sub>21</sub> and no poloxamer) | 63000 |
|    | 16 µg/mL rBPI <sub>21</sub> with 0.1% PLURONIC P123              | < 100 |
|    | 16 ug/mL rBPI <sub>21</sub> with 0.1% heat-treated PLURONIC P123 | 100   |
|    | 0.1% PLURONIC P123 control                                       | 70000 |
| 15 | 0.1% heat-treated PLURONIC P123 control                          | 70000 |
|    | 16 µg/mL rBPI <sub>21</sub> with 0.1% heat-treated PLURONIC P104 | 100   |

007201" 50296960



## EXAMPLE 2

### BACTERICIDAL ACTIVITY OF COMPOSITIONS CONTAINING BPI PROTEIN PRODUCT AND POLOXAMER 333 ON *S. AUREUS* AND *A. BAUMANNII* IN SERUM, BROTH OR WATER

5           The bactericidal activity of therapeutic compositions comprising BPI  
protein product and either poloxamer 333 (PLURONIC P103, BASF Wyandotte  
Corp.) or heat-treated PLURONIC P103, was evaluated against the clinical  
isolates of Example 1. Therapeutic compositions comprising 160  $\mu\text{g/mL}$  rBPI<sub>21</sub>  
and varying concentrations of either PLURONIC P103 or heat-treated PLURONIC  
10 P103 were formulated by diluting a 2 mg/mL solution of "non-formulated" rBPI<sub>21</sub>  
(in buffer comprising 5 mM sodium citrate and 150 mM NaCl, without any  
surfactants) with the appropriate amounts of PLURONIC P103 or heat-treated  
PLURONIC P103 solutions. A "formulated" rBPI<sub>21</sub> solution containing 2 mg/mL  
rBPI<sub>21</sub>, 0.2% poloxamer 188 (PLURONIC F68, BASF Wyandotte Corp.), 0.002%  
15 TWEEN 80 (polysorbate 80, ICI Americas, Wilmington, DE), 5 mM sodium  
citrate and 150 mM NaCl was also tested for comparison. Poloxamer control  
solutions containing only 0.1% PLURONIC P103 or 0.1% heat-treated  
PLURONIC P103, and no rBPI<sub>21</sub>, were also prepared.

          A 0.1% (w/v) solution of PLURONIC P103 was prepared by  
20 stirring the PLURONIC P103 in deionized water until dissolved and filtering the  
solution through a 0.22 $\mu\text{m}$  cellulose acetate polystyrene filter unit (Corning Inc.,  
Corning, NY). Sterile stock solutions of heat-treated PLURONIC P103 were  
prepared using the following procedure: (1) making a 0.1% (w/v) solution of  
PLURONIC P103 in deionized water, (2) boiling the solution for 30 minutes, (3)  
25 allowing it to cool to room temperature, (4) stirring until the PLURONIC P103  
was completely solubilized, and (5) filtering the solution through a 0.22 $\mu\text{m}$   
Acrodisc filter unit (Gelman Sciences, Ann Arbor, MI) for sterilization.

          The bacteria to be used in the assays were grown on tryptic soy  
agar (TSA) plates (Remel, Catalog #01-920, Lenexa, KN) for 24 hours. The *S.*  
30 *aureus* were grown for an additional 2 hours in Fildes enriched medium. A

09696709 102400

bacterial stock emulsion was prepared by emulsifying bacterial colonies in sterile deionized water to approximately  $2.2$  to  $3.8 \times 10^8$  colony forming units (CFU)/mL as measured by a Microscan® Turbidity Meter (Dade Microscan, West Sacramento, CA), and diluting further by 1:10 in water. Assays were conducted in 96-well flat-bottom microtiter plates (Corning, catalog# 25860-96) by adding to each well:  $170 \mu\text{L}$  of serum (Sigma #S1764, St. Louis, MO), tryptic soy broth (TSB, Remel, catalog #08-942, Lenexa, KN) or sterile water for injection (Kendall McGaw);  $10 \mu\text{L}$  of the bacterial emulsion (or water, as a control);  $20 \mu\text{L}$  of the indicated  $160 \mu\text{g/mL}$  rBPI<sub>21</sub>/poloxamer therapeutic composition (or the poloxamer control solution or water alone as a control). The final concentrations of bacteria in each well were about  $4$  to  $7 \times 10^5$  CFU/mL. The well contents were mixed and the plates were incubated at  $37^\circ\text{C}$  for 4 hours. Following incubation, the remaining colony forming units (CFU) in each well were counted at a  $10^{-2}$  dilution by plating  $10 \mu\text{L}$  from each well onto TSA plates. The TSA plates were incubated at  $37^\circ\text{C}$  for 24 hours and the number of bacterial colonies were visually counted. Results are shown below in Table 3; colony counts for the control wells are shown below in Tables 4 and 5.

09696709-102400

Table 3

| R<br>o<br>w<br>N<br>o. | Contents of<br>well  | 100's of CFU remaining after<br>incubation with serum and<br>16 µg/mL rBPI <sub>21</sub> formulated<br>with poloxamer at: |                                     |                                     |                                      | 100's of CFU remaining after<br>incubation with broth and<br>16 µg/mL rBPI <sub>21</sub> formulated<br>with poloxamer at: |                                     |                                     |                                      | 100's of CFU remaining after<br>incubation with water and<br>16 µg/mL rBPI <sub>21</sub> formulated<br>with poloxamer at: |                                     |                                     |                                      |
|------------------------|--|---|-------------------------------------|-------------------------------------|--------------------------------------|---|-------------------------------------|-------------------------------------|--------------------------------------|---|-------------------------------------|-------------------------------------|--------------------------------------|
|                        |  | 0.1 %<br>Formu-<br>lation<br>Conc.  | 0.05 %<br>Formu-<br>lation<br>Conc. | 0.01 %<br>Formu-<br>lation<br>Conc. | 0.005 %<br>Formu-<br>lation<br>Conc. | 0.1 %<br>Formu-<br>lation<br>Conc.  | 0.05 %<br>Formu-<br>lation<br>Conc. | 0.01 %<br>Formu-<br>lation<br>Conc. | 0.005 %<br>Formu-<br>lation<br>Conc. | 0.1 %<br>Formu-<br>lation<br>Conc.  | 0.05 %<br>Formu-<br>lation<br>Conc. | 0.01 %<br>Formu-<br>lation<br>Conc. | 0.005 %<br>Formu-<br>lation<br>Conc. |
|                        | (starting rBPI <sub>21</sub><br>solution;<br>type of<br>poloxamer<br>preparation;<br>organism) |   |                                     |                                     |                                      |   |                                     |                                     |                                      |   |                                     |                                     |                                      |
| A                      | NF rBPI <sub>21</sub> +<br>heat-treated<br>P103 +<br><i>A. baumannii</i>                       | > 2000  | > 2000                              | > 2000                              | > 2000                               | 0   | 0                                   | 0                                   | 0                                    | 0   | 0                                   | 0                                   | 0                                    |
| B                      | NF rBPI <sub>21</sub> +<br>P103 +<br><i>A. baumannii</i>                                       | > 2000  | > 2000                              | > 2000                              | > 2000                               | 0   | 0                                   | 0                                   | 0                                    | 0   | 0                                   | 0                                   | 0                                    |
| D                      | F rBPI <sub>21</sub> +<br>heat-treated<br>P103 +<br><i>A. baumannii</i>                        | > 2000  | > 2000                              | > 2000                              | > 2000                               | 0   | 0                                   | *                                   | 0                                    | 0   | > 2000                              | 0                                   | 0                                    |

- 27 -

Table 3

| R<br>o<br>w<br>N<br>o. | Contents of<br>well   | 100's of CFU remaining after<br>incubation with serum and<br>16 µg/mL rBPI <sub>21</sub> formulated<br>with poloxamer at: |                                     |                                     |                                      | 100's of CFU remaining after<br>incubation with broth and<br>16 µg/mL rBPI <sub>21</sub> formulated<br>with poloxamer at: |                                     |                                     |                                      | 100's of CFU remaining after<br>incubation with water and<br>16 µg/mL rBPI <sub>21</sub> formulated<br>with poloxamer at: |                                     |                                     |                                      |
|------------------------|---|---|-------------------------------------|-------------------------------------|--------------------------------------|---|-------------------------------------|-------------------------------------|--------------------------------------|---|-------------------------------------|-------------------------------------|--------------------------------------|
|                        |   | 0.1 %<br>Formu-<br>lation<br>Conc.  | 0.05 %<br>Formu-<br>lation<br>Conc. | 0.01 %<br>Formu-<br>lation<br>Conc. | 0.005 %<br>Formu-<br>lation<br>Conc. | 0.1 %<br>Formu-<br>lation<br>Conc.  | 0.05 %<br>Formu-<br>lation<br>Conc. | 0.01 %<br>Formu-<br>lation<br>Conc. | 0.005 %<br>Formu-<br>lation<br>Conc. | 0.1 %<br>Formu-<br>lation<br>Conc.  | 0.05 %<br>Formu-<br>lation<br>Conc. | 0.01 %<br>Formu-<br>lation<br>Conc. | 0.005 %<br>Formu-<br>lation<br>Conc. |
| E                      | F rBPI <sub>21</sub> +<br>P103 +<br><i>A. baumannii</i>               | > 2000  | > 2000                              | > 2000                              | > 2000                               | 0   | 0                                   | 51                                  | 252                                  | 0   | 0                                   | 0                                   | 0                                    |
| G                      | NF rBPI <sub>21</sub> +<br>heat-treated<br>P103 +<br><i>S. aureus</i> | > 1000  | > 1000                              | > 1000                              | > 1000                               | > 2000  | > 2000                              | > 2000                              | > 2000                               | 0   | 0                                   | 0                                   | 0                                    |

NF = non-formulated, i.e. prepared without surfactants

F = formulated with 0.2 % poloxamer 188 and 0.002 % polysorbate 80

\* = Contaminated

**Table 4**  
Growth Controls for *A. baumannii* (in 100's of CFUs)

|       |                                 |         |
|-------|---------------------------------|---------|
| Serum | NF rBPI <sub>21</sub> (no P103) | > 2000  |
|       | bacteria only                   | > 2000  |
|       | 0.1% heat-treated P103 (no BPI) | > 2000* |
|       | 0.1% P103 (no BPI)              | > 2000  |
| Broth | NF rBPI <sub>21</sub> (no P103) | > 5000  |
|       | bacteria only                   | > 5000  |
|       | 0.1% heat-treated P103 (no BPI) | > 5000  |
|       | 0.1% P103 (no BPI)              | > 5000  |
| Water | NF rBPI <sub>21</sub>           | 519     |
|       | bacteria only                   | > 2000  |
|       | 0.1% heat-treated P103 (no BPI) | > 2000  |
|       | 0.1% P103 (no BPI)              | > 2000  |

\*Contaminated

NF=non-formulated, i.e., prepared without surfactants

**Table 5**  
Growth controls for *S. aureus* (in 100's of CFUs)

| Serum and <i>S. aureus</i> | Serum and <i>S. aureus</i> and 0.1% heat-treated P103 | Broth and <i>S. aureus</i> | Broth and <i>S. aureus</i> and 0.1% heat-treated P103 | Water and <i>S. aureus</i> | Water and <i>S. aureus</i> and 0.1% heat-treated P103 |
|----------------------------|---|----------------------------|---|----------------------------|---|
| 2260                       | 2540  | 2960                       | 4240  | 550                        | 390   |

Additional experiments were performed to test therapeutic compositions, prepared by diluting a variety of formulated BPI protein products with heat-treated PLURONIC P104 solution, and tested against *A. baumannii* in serial 2-fold dilutions of serum. In these experiments, it was noted that some bactericidal activity was observed at lower serum concentrations (as evidenced by a serial 50% reduction in CFUs that correlated to the serial 2-fold reduction in serum concentration). For rBPI<sub>23</sub>, bactericidal activity was observed at serum concentrations of 12.5% and lower. For rBPI<sub>21</sub>, bactericidal activity was observed at serum concentrations of 6.25% and lower. For rBPI<sub>42</sub> dimer and rBPI<sub>50</sub>, bactericidal activity was observed at dilutions of 1.6% and lower.

In other experiments performed in a similar manner with Microscan® Pluronic Inoculum Water (Dade Microscan, West Sacramento, CA), this product exhibited bactericidal activity enhancing effect. In preliminary experiments performed in a similar manner with poloxamer 335 (PLURONIC P105, BASF Wyendotte Corp.), this poloxamer was also observed to have some bactericidal activity enhancing effect.

In further experiments, the bactericidal activity of therapeutic compositions comprising BPI protein product and a poloxamer surfactant was evaluated against clinical isolates of *Neisseria meningitidis* (Type C) (Microscan® ID No. 410-001), *Pseudomonas aeruginosa* (strain 12.4.4, provided by S.M. Opal, Brown University, Providence, Rhode Island; referenced in Ammons *et al.*, *J. Infect. Diseases*, 170:1473-82 (1994)), and *Acinetobacter baumannii* (Microscan® ID No. 12300). The following therapeutic compositions were prepared, comprising 2 mg/mL rBPI<sub>21</sub>; 0.2% of either (a) poloxamer 188 (PLURONIC F68), (b) poloxamer 333 (PLURONIC P103), (c) poloxamer 334 (PLURONIC P104), (d) poloxamer 335 (PLURONIC P105) or (e) poloxamer 403 (PLURONIC P123); 0.002% polysorbate 80 (TWEEN 80); 5mM sodium citrate; and 150 mM NaCl.

Poloxamer control solutions containing only 0.2% PLURONIC P123, P103 or F68, and no rBPI<sub>21</sub>, were also prepared.

5 The bacteria to be used in these additional assays were grown for approximately 24 hours on tryptic soy agar (TSA) plates (Remel, Catalog #01-920, Lenexa, KN) for *P. aeruginosa* or *A. baumannii* and chocolate agar plates (Remel Catalog # 01-301, Lenexa, KN) for *N. meningitidis*. A bacterial stock emulsion was prepared by emulsifying bacterial colonies in sterile saline (0.9% sodium chloride Irrigation water, Kendall McGaw Laboratory, Irvine, CA) to an equivalent of a 0.5 McFarland standard as measured by a  
10 Microscan® Turbidity Meter (Dade Microscan, West Sacramento, CA), and diluting further by 1:10 in saline. Assays were conducted in a final volume of 1 mL by adding 982 or 974 µL of Mueller-Hinton Broth with 2% Fildes Enrichment (Remel, Catalog #06-1496, Lenexa, KN) for *N. meningitidis* or of Mueller-Hinton Broth plus Cations (CSMHB, Remel) for *P. aeruginosa* to 4.5  
15 mL polypropylene tubes (Nalgene Cryovial, Nalge Co., Rochester, NY), followed by 10 µL of the bacterial emulsion (or broth media, as a control); and 8 or 16 µL of the 2 mg/mL rBPI<sub>21</sub>/poloxamer therapeutic composition. The tubes were mixed by vortexing and incubated at 37°C for 8 hours. Following incubation, the remaining colony forming units (CFU) were counted at varying  
20 dilutions ( $10^{-2}$  to  $10^{-7}$ ) by plating 10 µl or 100 µl of an appropriate dilution onto chocolate agar or TSA plates. The chocolate agar or TSA plates were incubated at 37°C (with 5% CO<sub>2</sub> for the *N. meningitidis* plates) for approximately 24 hours and the number of bacterial colonies were visually counted. Results are shown below in Tables 6 and 7.

004201" 50796960

Table 6

| <i>N. meningitidis</i> <sup>a</sup>                             | CFU                  |
|---|----------------------|
| Control   | 9.5x10 <sup>7</sup>  |
| 0.2% PLURONIC P123 Control <sup>b</sup>                         | 7.8x10 <sup>7</sup>  |
| 16μg/mL rBPI <sub>21</sub> with 0.2% PLURONIC P103 <sup>b</sup> | 3x10 <sup>3</sup>    |
| 32μg/mL rBPI <sub>21</sub> with 0.2% PLURONIC P103 <sup>b</sup> | 3x10 <sup>3</sup>    |
| 0.2% PLURONIC F68 Control <sup>b</sup>                          | 10.1x10 <sup>7</sup> |
| 16μg/mL rBPI <sub>21</sub> with 0.2% PLURONIC F68 <sup>b</sup>  | 4.22x10 <sup>6</sup> |
| 32μg/mL rBPI <sub>21</sub> with 0.2% PLURONIC F68 <sup>b</sup>  | 1.2x10 <sup>3</sup>  |

10    <sup>a</sup> At t = 0, there were 2.02x10<sup>5</sup> organisms

<sup>b</sup> Also contains 0.002% TWEEN 80 (polysorbate 80)

004207 60296960



Table 7

| <i>P. aeruginosa</i> <sup>a</sup>                   |  | CFU                            |
|---|--|--------------------------------|
| Media Control                                       |  | 6.0 x 10 <sup>7</sup>          |
| 32 µg/ml rBPI <sub>21</sub> with 0.2% PLURONIC F68  |  | 1.2 x 10 <sup>8</sup>          |
| 32 µg/ml rBPI <sub>21</sub> with 0.2% PLURONIC P103 |  | < 10 <sup>6</sup> <sup>b</sup> |
| 32 µg/ml rBPI <sub>21</sub> with 0.2% PLURONIC P104 |  | 3 x 10 <sup>7</sup>            |
| 32 µg/ml rBPI <sub>21</sub> with 0.2% PLURONIC P105 |  | < 10 <sup>6</sup> <sup>b</sup> |
| 32 µg/ml rBPI <sub>21</sub> with 0.2% PLURONIC P123 |  | < 10 <sup>6</sup> <sup>b</sup> |
|   |  |                                |
| <i>A. baumannii</i> <sup>c</sup>                    |  | CFU                            |
| Media Control                                       |  | 1.06 x 10 <sup>7</sup>         |
| 16 µg/ml rBPI <sub>21</sub> with 0.2% PLURONIC F68  |  | 2.43 x 10 <sup>7</sup>         |
| 16 µg/ml rBPI <sub>21</sub> with 0.2% PLURONIC P103 |  | < 10 <sup>d</sup>              |
| 16 µg/ml rBPI <sub>21</sub> with 0.2% PLURONIC P104 |  | < 10 <sup>d</sup>              |
| 16 µg/ml rBPI <sub>21</sub> with 0.2% PLURONIC P105 |  | < 10 <sup>d</sup>              |
| 16 µg/ml rBPI <sub>21</sub> with 0.2% PLURONIC P123 |  | 2.7 x 10 <sup>2</sup>          |

<sup>a</sup> At t=0, there were 6.4 x 10<sup>5</sup> CFUs

<sup>b</sup> No CFUs at tested dilutions of 10<sup>-6</sup> and 10<sup>-7</sup>

<sup>c</sup> At t=0, there were 4.7 x 10<sup>4</sup> CFUs

<sup>d</sup> No CFUs at tested dilutions of 10<sup>-1</sup> and 10<sup>-2</sup>

004201" 60296960

### EXAMPLE 3

#### BACTERICIDAL ACTIVITY OF COMPOSITIONS CONTAINING BPI PROTEIN PRODUCT AND POLOXAMER 333 OR POLOXAMER 334 ON A VARIETY OF BACTERIA IN WATER

- 5           The bactericidal activity of therapeutic compositions comprising BPI protein product and heat-treated PLURONIC P103 or heat-treated PLURONIC P104, was evaluated against the *S. aureus* and *A. baumannii* clinical isolates of Example 1 and the additional organisms *S. pneumoniae* (Microscan® ID no. 145) and *E. faecium* (Microscan® ID no. 15773).
- 10   Therapeutic compositions comprising 500 µg/mL rBPI<sub>21</sub> in a 0.075% (w/v) concentration of either heat-treated PLURONIC P103 or heat-treated PLURONIC P104 were formulated by diluting a 2 mg/mL solution of "non-
- 15   formulated" rBPI<sub>21</sub> or "formulated" rBPI<sub>21</sub> with the appropriate amounts of 0.1% heat-treated PLURONIC P103 or heat-treated PLURONIC P104 solutions. Compositions comprising 500 µg/mL non-formulated rBPI<sub>21</sub> in water alone (without any poloxamers) and poloxamer control solutions containing only 0.1% heat-treated P103 or heat-treated P104 (and no rBPI<sub>21</sub>) were also prepared. A "formulated" rBPI<sub>23</sub> therapeutic composition containing 1 mg/mL rBPI<sub>23</sub>, 0.1% PLURONIC F68 and 0.002% TWEEN 80 was also
- 20   tested for comparison.

- Sterile stock solutions of heat-treated PLURONIC P103 or heat-treated PLURONIC P104 were prepared using the following procedure: (1) making a 0.1% (w/v) solution of the poloxamer in deionized water, (2) heating the solution to a boil, (3) allowing it to cool to room temperature, (4) stirring
- 25   until the PLURONIC P103 was completely solubilized, and (5) filtering the solution through a 0.22µm Nalgene filter for sterilization.

- The *S. aureus*, *E. faecium* and *A. baumannii* bacteria were grown on TSA plates (Remel, Catalog #01-920, Lenexa, KN), and the *S. pneumoniae* were grown on 5% sheep blood agar plates (Remel, Catalog# 01-
- 30   200, Lenexa, KN) for 24 hours. A bacterial stock emulsion was prepared by

004207" 60296960

emulsifying bacterial colonies in sterile deionized water to approximately 2.2 to  $3.8 \times 10^8$  CFU/mL as measured by a Microscan® Turbidity Meter, and diluting further by 1:10 in water. Assays for rBPI<sub>21</sub> therapeutic compositions were conducted in 96-well flat-bottom microtiter plates (Corning, catalog#

- 5 25860-96) by adding to each well: 185  $\mu$ L of TSB (Remel, catalog #08-942, Lenexa, KN) or sterile water for injection (Kendall McGaw); 8  $\mu$ L of the bacterial emulsion; 6.3  $\mu$ L of the indicated 500  $\mu$ g/mL rBPI<sub>21</sub>/poloxamer therapeutic composition (or poloxamer control solution or water alone). The final concentrations of bacteria in each well were about 4 to  $7 \times 10^5$  CFU/mL.
- 10 Assays for the rBPI<sub>23</sub> therapeutic composition were conducted in the same way, except 178  $\mu$ L of broth or water and 13  $\mu$ L of the 500  $\mu$ g/mL rBPI<sub>23</sub> composition were added. The well contents were mixed and the plates were incubated at 37°C. The CFUs in each well were counted at  $10^{-2}$  and  $10^{-4}$
- 15 and 3 hours, respectively, are shown below in Tables 8 and 9.

In a preliminary experiment using therapeutic compositions containing rBPI<sub>21</sub> and heat-treated PLURONIC P104, it was noted that adding the therapeutic composition immediately after the diluent (*e.g.* water), before addition of the bacteria, provided greater enhancement of the bactericidal

20 activity of rBPI<sub>21</sub> compared to adding the same therapeutic composition after adding bacteria. In another preliminary experiment performed using the same gram-positive and gram-negative organisms, with therapeutic compositions prepared by diluting non-formulated rBPI<sub>21</sub> with PLURONIC P103 and PLURONIC P104 solutions, no bactericidal activity was observed against the

25 gram-positive organisms in broth at concentrations of up to 64  $\mu$ g/mL of the rBPI<sub>21</sub> therapeutic compositions.

09696709-102400

Table 8: Incubation for 30 minutes

|                            |       | NF<br>rBPL <sub>21</sub><br>alone | NF<br>rBPL <sub>21</sub><br>with<br>0.075%<br>heat-<br>treated<br>P103 | NF<br>rBPL <sub>21</sub><br>with<br>0.075%<br>heat-<br>treated<br>P104 | F<br>rBPL <sub>21</sub><br>alone | Con-<br>trol | F<br>rBPL <sub>23</sub><br>alone | F<br>rBPL <sub>23</sub><br>with<br>0.075%<br>heat-<br>treated<br>P103 | F<br>rBPL <sub>23</sub><br>with<br>0.075%<br>heat-<br>treated<br>P104 | Con-<br>trol |     |
|----------------------------|-------|-----------------------------------|--|--|----------------------------------|--------------|----------------------------------|---|---|--------------|-----|
| <i>S. pneumo-<br/>niae</i> | water | 100<br>CFUs                       | 61   | 47   | 58                               | 57           | 75                               | 66  | 58  | 43           | 47  |
|                            | water | 10000<br>CFUs                     | 0  | 0  | 1                                | 0            | 1                                | 1   | 1   | 0            | 0   |
|                            | broth | 100<br>CFUs                       | 290  | 305  | 224                              | 355          | 389                              | 337   | 340   | 350          | 350 |
|                            | broth | 10000<br>CFUs                     | 4  | 3  | 0                                | 4            | 4                                | 4   | 5   | 7            | 1   |

Table 8: Incubation for 30 minutes

|                  |       | NF<br>rBPL <sub>21</sub><br>alone | NF<br>rBPL <sub>21</sub><br>with<br>0.075 %<br>heat-<br>treated<br>P103 | NF<br>rBPL <sub>21</sub><br>with<br>0.075 %<br>heat-<br>treated<br>P104 | F<br>rBPL <sub>21</sub><br>alone | Con-<br>trol | F<br>rBPL <sub>23</sub><br>alone | F<br>rBPL <sub>23</sub><br>with<br>0.075 %<br>heat-<br>treated<br>P103 | F<br>rBPL <sub>23</sub><br>with<br>0.075 %<br>heat-<br>treated<br>P104 | Con-<br>trol |
|------------------|-------|-----------------------------------|---|---|----------------------------------|--------------|----------------------------------|--|--|--------------|
| <i>S. aureus</i> | water | 100<br>CFUs                       | 315   | 227   | 305                              | TNTC         | TNTC                             | TNTC   | 398  | TNTC         |
|                  | water | 10000<br>CFUs                     | 2   | 1   | 5                                | 36           | 18                               | 3  | 1  | 63           |
|                  | broth | 100<br>CFUs                       | TNTC  | TNTC  | TNTC                             | TNTC         | TNTC                             | TNTC   | TNTC   | TNTC         |
|                  | broth | 10000<br>CFUs                     | 57  | 68  | 49                               | 75           | 60                               | 54   | 75   | 59           |

**Table 8: Incubation for 30 minutes**

|                   |       | NF<br>rBPI <sub>21</sub><br>alone | NF<br>rBPI <sub>21</sub><br>with<br>0.075 %<br>heat-<br>treated<br>P103 | NF<br>rBPI <sub>21</sub><br>with<br>0.075 %<br>heat-<br>treated<br>P104 | F<br>rBPI <sub>21</sub><br>alone | Con-<br>trol | F<br>rBPI <sub>23</sub><br>alone | F<br>rBPI <sub>23</sub><br>with<br>0.075 %<br>heat-<br>treated<br>P103 | F<br>rBPI <sub>23</sub><br>with<br>0.075 %<br>heat-<br>treated<br>P104 | Con-<br>trol |
|-------------------|-------|-----------------------------------|---|---|----------------------------------|--------------|----------------------------------|--|--|--------------|
| <i>E. faecium</i> | water | 100<br>CFUs                       | 50  | 33  | 122                              | 396          | TNTC                             | 180  | 165  | TNTC         |
|                   | water | 10000<br>CFUs                     | 1   | 0   | 3                                | 7            | 15                               | 3  | 4  | 35           |
|                   | broth | 100<br>CFUs                       | TNTC  | TNTC  | TNTC                             | TNTC         | TNTC                             | TNTC   | TNTC   | TNTC         |
|                   | broth | 10000<br>CFUs                     | 89  | 28  | 50                               | 55           | 57                               | 51   | 39   | 38           |

Table 8: Incubation for 30 minutes

|                              |       | NF<br>rBPI <sub>21</sub><br>alone | NF<br>rBPI <sub>21</sub><br>with<br>0.075%<br>heat-<br>treated<br>P103 | NF<br>rBPI <sub>21</sub><br>with<br>0.075%<br>heat-<br>treated<br>P104 | F<br>rBPI <sub>21</sub><br>alone | Con-<br>trol | F<br>rBPI <sub>23</sub><br>alone | F<br>rBPI <sub>23</sub><br>with<br>0.075%<br>heat-<br>treated<br>P103 | F<br>rBPI <sub>23</sub><br>with<br>0.075%<br>heat-<br>treated<br>P104 | Con-<br>trol |
|------------------------------|-------|-----------------------------------|--|--|----------------------------------|--------------|----------------------------------|---|---|--------------|
| A.<br><i>anitra-<br/>tus</i> | water | 100<br>CFUs                       | 73   | 0  | 49                               | TNTC         | 203                              | 0   | 0   | TNTC         |
|                              | water | 10000<br>CFUs                     | 0  | 0  | 1                                | 16           | 3                                | 0   | 0   | 17           |
|                              | broth | 100<br>CFUs                       | TNTC   | 68   | TNTC                             | TNTC         | TNTC                             | 33  | 67  | TNTC         |
|                              | broth | 10000<br>CFUs                     | 24   | 2  | 28                               | 44           | 29                               | 0   | 3   | 41           |

Table 9: Incubation for 3 hours

|                      |       | NF<br>rBPI <sub>21</sub><br>alone | NF rBPI <sub>21</sub><br>with<br>0.075 %<br>heat-<br>treated<br>P103 | NF rBPI <sub>21</sub><br>with<br>0.075 %<br>heat-treated<br>P104 | F rBPI <sub>21</sub><br>alone | Control | F rBPI <sub>23</sub><br>alone | F rBPI <sub>23</sub><br>with<br>0.075 %<br>heat-<br>treated<br>P103 | F rBPI <sub>23</sub><br>with<br>0.075 %<br>heat-<br>treated<br>P104 |
|----------------------|-------|-----------------------------------|--|--|-------------------------------|---------|-------------------------------|---|---|
| <i>S. pneumoniae</i> | water | 0                                 | 0  | 0  | 0                             | 0       | 0                             | 0   | 0   |
|                      | water |                                   |  |  |                               | 0       |                               |   |   |
|                      | broth | 447                               | 377  | 393  | 400                           | 337     | 360                           | 274   | 400   |
|                      | broth |                                   |  |  |                               | 0       |                               |   |   |
| <i>S. aureus</i>     | water | 12                                | 2  | 5  | 340                           | TNTC    | 840                           | 10  | 14  |
|                      | water |                                   |  |  |                               | 36      |                               |   |   |
|                      | broth | TNTC                              | TNTC   | TNTC   | TNTC                          | TNTC    | TNTC                          | TNTC  | TNTC  |
|                      | broth |                                   |  |  |                               | 144     |                               |   |   |



- 40 -

Table 9: Incubation for 3 hours

|                   |       | NF rBPI <sub>21</sub> alone | NF rBPI <sub>21</sub> with 0.075 % heat-treated P103 | NF rBPI <sub>21</sub> with 0.075 % heat-treated P104 | F rBPI <sub>21</sub> alone | Control | F rBPI <sub>23</sub> alone | F rBPI <sub>23</sub> with 0.075 % heat-treated P103 | F rBPI <sub>23</sub> with 0.075 % heat-treated P104 |
|-------------------|-------|-----------------------------|--|--|----------------------------|---------|----------------------------|---|---|
| <i>E. faecium</i> | water | 100 CFUs                    | 1  | 1  | 3                          | 28      | 498                        | 8   | 7   |
|                   | water | 10000 CFUs                  |  |  |                            | 36      |                            |   |   |
|                   | broth | 100 CFUs                    | TNTC   | TNTC   | TNTC                       | TNTC    | TNTC                       | TNTC  | TNTC  |
|                   | broth | 10000 CFUs                  |  |  |                            | 167     |                            |   |   |

Table 9: Incubation for 3 hours

|                     |       | NF<br>rBPI <sub>21</sub><br>alone | NF rBPI <sub>21</sub><br>with<br>0.075 %<br>heat-<br>treated<br>P103 | NF rBPI <sub>21</sub><br>with<br>0.075 %<br>heat-treated<br>P104 | F rBPI <sub>21</sub><br>alone | Control | F rBPI <sub>23</sub><br>alone | F rBPI <sub>23</sub><br>with<br>0.075 %<br>heat-<br>treated<br>P103 | F rBPI <sub>23</sub><br>with<br>0.075 %<br>heat-<br>treated<br>P104 |
|---------------------|-------|-----------------------------------|--|--|-------------------------------|---------|-------------------------------|---|---|
| <i>A. baumannii</i> | water | 100<br>CFUs                       | 0  | 0  | 0                             | TNTC    | 0                             | 0   | 0   |
|                     | water | 10000<br>CFUs                     |  |  |                               | 15      |                               |   |   |
|                     | broth | 100<br>CFUs                       | 58   | 0  | 27                            | TNTC    | 15                            | 0   | 0   |
|                     | broth | 10000<br>CFUs                     |  |  |                               | 263     |                               |   |   |

#### EXAMPLE 4

##### BACTERICIDAL ACTIVITY OF COMPOSITIONS CONTAINING BPI PROTEIN PRODUCT AND OTHER POLOXAMER SURFACTANTS

Therapeutic compositions comprising BPI protein product and  
5 other poloxamer surfactants, including poloxamer 101, poloxamer 105,  
poloxamer 108, poloxamer 122, poloxamer 123, poloxamer 124, poloxamer  
181, poloxamer 182, poloxamer 183, poloxamer 184, poloxamer 185,  
poloxamer 188, poloxamer 212, poloxamer 215, poloxamer 217, poloxamer  
231, poloxamer 234, poloxamer 235, poloxamer 237, poloxamer 238,  
10 poloxamer 282, poloxamer 284, poloxamer 288, poloxamer 331, poloxamer  
333, poloxamer 334, poloxamer 335, poloxamer 338, poloxamer 401,  
poloxamer 402, poloxamer 403, or poloxamer 407 [see, *e.g.*, *CTFA  
International Cosmetic Ingredient Dictionary*, Cosmetic, Toiletry and  
Fragrance Association, Inc., Washington, DC (1991)], especially at pages 447-  
15 451] are prepared and tested for capacity to enhance bactericidal activity of  
BPI protein products as described above in Examples 1, 2 and 3.

#### EXAMPLE 5

BACTERICIDAL ACTIVITY OF COMPOSITIONS CONTAINING  
BPI PROTEIN PRODUCT FORMULATED WITH POLOXAMER, WITH  
20 OR WITHOUT EDTA, IN SERUM, MUELLER-HINTON BROTH,  
TRYPTIC SOY BROTH, OR WATER

The bactericidal activity of therapeutic compositions comprising  
BPI protein product and PLURONIC F68, P103, P104, P105 or P123 were  
evaluated against the *S. aureus* and *A. baumannii* organisms of Example 1, the  
25 *S. pneumoniae* organism of Example 3, an *E. faecium* organism (Microscan®  
ID No. 16866), and a strain of *P. aeruginosa* from the American Type Culture  
Collection (ATCC No. 19660). Therapeutic compositions were formulated by  
adding the appropriate amount of poloxamer to a stock solution of 2.2 mg/mL  
rBPI<sub>21</sub> (5 mM sodium citrate, 150 mM NaCl, without poloxamer), to achieve  
30 the desired 0.2% (w/v) poloxamer concentration, followed by sterile filtration.

09696709-102400

Formulated product was stored at 2-8°C for up to 6 months. Sterile stock solutions of poloxamer were made by dissolving the poloxamer paste in water for injection (WFI, Kendall-McGaw) with mixing to a 1-5% concentration (w/v) at room temperature, followed by sterile filtration. Assays were

5 conducted in 96-well microtiter plates using WFI, tryptic soy broth (TSB, Remel, Lenexa, KN), Mueller-Hinton Broth plus Cations (CSMHB, Remel), or 40% pooled human serum in CSMHB (Sigma, St. Louis, MO) as growth media, according to the general procedure described above in Examples 2 and 3. The results (in colony forming units after 24 hours of incubation) are

10 displayed below in Table 10, and confirm that the poloxamers can enhance the bactericidal activity of BPI protein product.

004207" 60296960

Table 10

| Organism             | Medium | Control             | rBPL <sub>21</sub><br>only | rBPL <sub>21</sub><br>with F68 | rBPL <sub>21</sub><br>with<br>P103 | rBPL <sub>21</sub><br>with<br>P104 | rBPL <sub>21</sub><br>with<br>P105 | rBPL <sub>21</sub><br>with<br>P123 |
|----------------------|--------|---------------------|----------------------------|--------------------------------|------------------------------------|------------------------------------|------------------------------------|------------------------------------|
| <i>A. baumannii</i>  | Water  | 2x10 <sup>6</sup>   | <100                       | <100                           | <100                               | <100                               | <100                               | <100                               |
|                      | TSB    | 3x10 <sup>6</sup>   | 6x10 <sup>2</sup>          | 3x10 <sup>2</sup>              | <100                               | <100                               | <100                               | <100                               |
|                      | CSMHB  | 2x10 <sup>6</sup>   | NT                         | NT                             | <100                               | 100                                | <100                               | 300                                |
|                      | Serum  | 2x10 <sup>5</sup>   | 2x10 <sup>5</sup>          | 2x10 <sup>5</sup>              | 2x10 <sup>5</sup>                  | 2x10 <sup>5</sup>                  | 2x10 <sup>5</sup>                  | 3x10 <sup>3</sup>                  |
| <i>S. aureus</i>     | Water  | 8.2x10 <sup>5</sup> | 3.2x10 <sup>4</sup>        | 3.6x10 <sup>5</sup>            | 2.3x10 <sup>4</sup>                | 3.0x10 <sup>4</sup>                | NT                                 | 2.7x10 <sup>4</sup>                |
|                      | TSB    | 5.4x10 <sup>5</sup> | 5.7x10 <sup>5</sup>        | 7.5x10 <sup>5</sup>            | 6.0x10 <sup>5</sup>                | 7.2x10 <sup>5</sup>                | NT                                 | NT                                 |
|                      | CSMHB  | NT                  | NT                         | NT                             | NT                                 | NT                                 | NT                                 | NT                                 |
|                      | Serum  | 4.2x10 <sup>5</sup> | >1x10 <sup>5</sup>         | >1x10 <sup>5</sup>             | >1x10 <sup>5</sup>                 | NT                                 | NT                                 | NT                                 |
| <i>S. pneumoniae</i> | Water  | 3.2x10 <sup>5</sup> | NT                         | >1x10 <sup>5</sup>             | <100                               | <100                               | 400                                | <100                               |
|                      | TSB    | 3x10 <sup>5</sup>   | 5x10 <sup>4</sup>          | 4x10 <sup>4</sup>              | <100                               | 5x10 <sup>4</sup>                  | 3x10 <sup>3</sup>                  | <100                               |
|                      | CSMHB  | 1x10 <sup>7</sup>   | NT                         | NT                             | 2x10 <sup>3</sup>                  | 9x10 <sup>2</sup>                  | 3x10 <sup>4</sup>                  | 8x10 <sup>3</sup>                  |
|                      | Serum  | 3x10 <sup>5</sup>   | NT                         | 2.9x10 <sup>5</sup>            | 6x10 <sup>4</sup>                  | 6x10 <sup>4</sup>                  | 6x10 <sup>4</sup>                  | 6x10 <sup>4</sup>                  |

Table 10

| Organism             | Medium         | Control           | rBPI <sub>21</sub><br>only | rBPI <sub>21</sub><br>with F68 | rBPI <sub>21</sub><br>with<br>P103 | rBPI <sub>21</sub><br>with<br>P104 | rBPI <sub>21</sub><br>with<br>P105 | rBPI <sub>21</sub><br>with<br>P123 |
|----------------------|----------------|-------------------|----------------------------|--------------------------------|------------------------------------|------------------------------------|------------------------------------|------------------------------------|
| <i>E. faecium</i>    | Water<br>TSB   | 4x10 <sup>5</sup> | 100                        | 3x10 <sup>3</sup>              | 100                                | 300                                | 300                                | <100                               |
|                      |                | 5x10 <sup>5</sup> | 5x10 <sup>5</sup>          | 5x10 <sup>5</sup>              | 4x10 <sup>3</sup>                  | 3.1x10 <sup>5</sup>                | 6x10 <sup>4</sup>                  | 1x10 <sup>3</sup>                  |
|                      | CSMHB<br>Serum | 1x10 <sup>7</sup> | NT                         | NT                             | 8x10 <sup>4</sup>                  | 4x10 <sup>5</sup>                  | 2x10 <sup>5</sup>                  | 6x10 <sup>5</sup>                  |
|                      |                | 1x10 <sup>8</sup> | NT                         | NT                             | 5x10 <sup>7</sup>                  | 7x10 <sup>7</sup>                  | 5x10 <sup>7</sup>                  | 1x10 <sup>8</sup>                  |
| <i>P. aeruginosa</i> | Water<br>TSB   | 1x10 <sup>7</sup> | NT                         | NT                             | 3x10 <sup>3</sup>                  | 2x10 <sup>3</sup>                  | 7x10 <sup>4</sup>                  | 2x10 <sup>3</sup>                  |
|                      |                | NT                | NT                         | NT                             | NT                                 | NT                                 | NT                                 | NT                                 |
|                      | CSMHB<br>Serum | 1x10 <sup>8</sup> | NT                         | 4x10 <sup>7</sup>              | 1x10 <sup>7</sup>                  | 5x10 <sup>7</sup>                  | 3x10 <sup>7</sup>                  | 5x10 <sup>7</sup>                  |
|                      |                | 4x10 <sup>7</sup> | NT                         | 3x10 <sup>7</sup>              | 3x10 <sup>6</sup>                  | 2x10 <sup>7</sup>                  | 3x10 <sup>7</sup>                  | 2x10 <sup>7</sup>                  |

In additional experiments, the bactericidal activity of therapeutic compositions comprising BPI protein product with a poloxamer surfactant and further comprising varying concentrations of EDTA were evaluated against *P. aeruginosa* (ATCC 19660). Therapeutic compositions were formulated as described above to achieve the desired concentrations of poloxamer and rBPI<sub>21</sub> in a buffer of 5mM sodium citrate, 150 mM NaCl and 0.002% polysorbate 80. Assays were conducted generally as described in Example 2 above for *P. aeruginosa* and *A. baumannii*. Results in colony forming units after approximately 24 hours of incubation are displayed below in Table 11, and show that the addition of EDTA can further enhance the bactericidal activity of BPI protein product formulated with PLURONIC P123.

004207" 60296960

Table 11

| CFU after incubation   |  |  |  |
|--|--|--|--|
| <i>P. aeruginosa</i><br>(ATCC No. 19660) <sup>a</sup>                                | 2 hours<br>incubation                      | 4 hours<br>incubation                      | 6 hours<br>incubation                      |
| Media Control (Mueller-Hinton<br>plus cations)                                       | 4.2x10 <sup>3</sup>                        | 1x10 <sup>5</sup>                          | 2.1x10 <sup>6</sup>                        |
| Placebo Control (Media with<br>formulation buffer and 0.05 %<br>EDTA)                | 1.3x10 <sup>3</sup>                        | 1.03x10 <sup>5</sup>                       | 5.4x10 <sup>6</sup>                        |
| 16μg/mL rBPI <sub>21</sub> with 0.2 %<br>PLURONIC P123 without EDTA                  | 7.0x10 <sup>3</sup><br>8.5x10 <sup>3</sup> | 4.5x10 <sup>4</sup><br>8.0x10 <sup>4</sup> | 5.4x10 <sup>5</sup><br>3.3x10 <sup>5</sup> |
| 16μg/mL rBPI <sub>21</sub> with 0.2 %<br>PLURONIC P123 + 0.05 %<br>EDTA <sup>b</sup> | 6.6x10 <sup>3</sup>                        | 1.34x10 <sup>5</sup>                       | 3.3x10 <sup>5</sup>                        |
| 128μg/mL rBPI <sub>21</sub> with 0.2 %<br>PLURONIC P123 without EDTA                 | 5.0x10 <sup>3</sup>                        | 3x10 <sup>4</sup>                          | 1x10 <sup>5</sup>                          |
| 128μg/mL rBPI <sub>21</sub> with 0.2 %<br>PLURONIC P123 + 0.05 %<br>EDTA             | 1.7x10 <sup>3</sup>                        | 3x10 <sup>3</sup>                          | 5x10 <sup>2</sup>                          |

a At t=0, there were 4.5 x 10<sup>3</sup> organisms.

b Also contains 0.002 % TWEEN 80 (polysorbate 80).

## EXAMPLE 6

### EFFECT OF COMPOSITIONS CONTAINING BPI PROTEIN PRODUCT AND POLOXAMER IN THE PRESENCE OR ABSENCE OF EDTA ON THE SUSCEPTIBILITY OF VARIOUS ORGANISMS TO ANTIBIOTICS

The effect of therapeutic compositions of rBPI<sub>21</sub> formulated with poloxamer, with or without EDTA, was evaluated on the antibiotic susceptibility of the multiple drug resistant *A. baumannii*, *S. pneumoniae*, *E. faecium* and *P. aeruginosa* organisms of Example 5. Therapeutic compositions



were prepared containing 2 mg/mL rBPI<sub>21</sub> (5 mM sodium citrate, 150 mM NaCl) with a 0.2% (w/v) concentration of PLURONIC F68, P103, P104, P105 or P123. The effect on the antibiotic susceptibility of the organisms was determined in Mueller-Hinton Broth plus Cations (CSMHB, Remel), or 40% pooled human serum in CSMHB (Sigma, St. Louis, MO), as follows.

Isolated colonies of the organism from overnight cultures were suspended in Microscan® Inoculum Water to a concentration equivalent to a 0.5 McFarland Standard (approximately  $1 \times 10^8$  CFU/ml), determined using a Microscan® turbidimeter. Aliquots were transferred to either CSMHB or 40% pooled human serum in CSMHB. Each tube contained either a final concentration of 16 µg/mL rBPI<sub>21</sub> or an equivalent volume of control buffer. Minimal inhibitory concentrations (MIC) for each antibiotic tested, *i.e.* the lowest concentration of antibiotic which inhibits visible growth, were determined using gram-negative (MB and MC) and gram-positive (MA) Sensititre Trays (Radiometer America, Westlake, OH), which allow for the rapid and simultaneous survey of a broad spectrum of standard antibiotics. Any other antimicrobial panel systems known in the art, such as the Microscan® (Dade Microscan, Sacramento, CA), Pasco (DIFCO, Detroit, MI) and Alamar (Alamar, Sacramento, CA) systems, may alternatively be used to assay for antibiotic susceptibility.

Tables 12-15 below display a summary of the results of the antibiotic screening panels, reported for each strain tested as the MIC of the tested antibiotics in the presence of the indicated rBPI<sub>21</sub> therapeutic composition. The antibiotic susceptibility standards (interpretation of an MIC as clinically resistant (R), intermediate (I) or susceptible (S) according to NCCLS standards) applicable to the organism tested appear in superscript next to the MIC. These results indicate that the improvement in therapeutic effectiveness of antibiotics that is seen with the addition of BPI protein product can be further enhanced by various poloxamer formulations.

004201" 60296960

- 49 -

Table 12

Effect of BPI protein product formulation on antibiotic susceptibility of *P. aeruginosa*

|                   |             | Minimum Inhibitory Concentration (μg/mL) |                             |                              |                              |                              |                              |  |
|-------------------|-------------|--|-----------------------------|------------------------------|------------------------------|------------------------------|------------------------------|--|
| Antibiotic Tested | Medium Used | Control (no BPI)                         | rBPL <sub>21</sub> with F68 | rBPL <sub>21</sub> with P103 | rBPL <sub>21</sub> with P104 | rBPL <sub>21</sub> with P105 | rBPL <sub>21</sub> with P123 |  |
| Ceftizoxime       | CSMHB       | > 128 <sup>R</sup>                       | 32 <sup>I</sup>             | 16 <sup>I</sup>              | 128 <sup>R</sup>             | 32 <sup>I</sup>              | 128 <sup>R</sup>             |  |
|                   | Serum       | 128 <sup>R</sup>                         | > 128 <sup>R</sup>          | 16 <sup>I</sup>              | 128 <sup>R</sup>             | 16 <sup>I</sup>              | 16 <sup>I</sup>              |  |
| Ceftriaxone       | CSMHB       | > 128 <sup>R</sup>                       | 32 <sup>I</sup>             | 8 <sup>S</sup>               | 128 <sup>R</sup>             | 32 <sup>I</sup>              | 128 <sup>R</sup>             |  |
|                   | Serum       | 128 <sup>R</sup>                         | > 128 <sup>R</sup>          | 16 <sup>I</sup>              | 128 <sup>R</sup>             | 16 <sup>I</sup>              | 32 <sup>I</sup>              |  |
| Chloramphenicol   | CSMHB       | > 32 <sup>R</sup>                        | > 32 <sup>R</sup>           | 16 <sup>I</sup>              | > 32 <sup>R</sup>            | 16 <sup>I</sup>              | 16 <sup>I</sup>              |  |
|                   | Serum       | > 32 <sup>R</sup>                        | > 32 <sup>R</sup>           | 16 <sup>I</sup>              | 16 <sup>I</sup>              | 16 <sup>I</sup>              | 16 <sup>I</sup>              |  |

- 50 -

Table 13

Effect of BPI protein product formulation on antibiotic susceptibility of *A. baumannii*

|                   |             | Minimum Inhibitory Concentration (μg/mL) |                             |                              |                              |                              |                              |  |
|-------------------|-------------|--|-----------------------------|------------------------------|------------------------------|------------------------------|------------------------------|--|
| Antibiotic Tested | Medium Used | Control (no BPI)                         | rBPI <sub>21</sub> with F68 | rBPI <sub>21</sub> with P103 | rBPI <sub>21</sub> with P104 | rBPI <sub>21</sub> with P105 | rBPI <sub>21</sub> with P123 |  |
| Ceftazidime       | CSMHB       | 16 <sup>I</sup>                          | 32 <sup>R</sup>             | <4 <sup>S</sup>              | <4 <sup>S</sup>              | <4 <sup>S</sup>              | <4 <sup>S</sup>              |  |
|                   | Serum       | >32 <sup>R</sup>                         | 32 <sup>R</sup>             | 16 <sup>I</sup>              | 16 <sup>I</sup>              | 16 <sup>I</sup>              | 16 <sup>I</sup>              |  |
| Ceftriaxone       | CSMHB       | 128 <sup>R</sup>                         | >128 <sup>R</sup>           | <1 <sup>S</sup>              | <1 <sup>S</sup>              | <1 <sup>S</sup>              | 4 <sup>S</sup>               |  |
|                   | Serum       | >128 <sup>R</sup>                        | >128 <sup>R</sup>           | >128 <sup>R</sup>            | >128 <sup>R</sup>            | >128 <sup>R</sup>            | >128 <sup>R</sup>            |  |
| Chloramphenicol   | CSMHB       | >4 <sup>R</sup>                          | 1 <sup>S</sup>              | <0.5 <sup>S</sup>            | 1 <sup>S</sup>               | <0.5 <sup>S</sup>            | <0.5 <sup>S</sup>            |  |
|                   | Serum       | >4 <sup>R</sup>                          | >4 <sup>R</sup>             | 2 <sup>S</sup>               | 4 <sup>R</sup>               | >4 <sup>R</sup>              | >4 <sup>R</sup>              |  |

- 51 -

Table 14

Effect of BPI protein product formulation on antibiotic susceptibility of *S. pneumoniae*

| Antibiotic Tested | Medium Used | Minimum Inhibitory Concentration ( $\mu\text{g/mL}$ ) |                             |                              |                              |                              |                              |
|-------------------|-------------|---|-----------------------------|------------------------------|------------------------------|------------------------------|------------------------------|
|                   |             | Control (no BPI)                                      | rBPI <sub>21</sub> with F68 | rBPI <sub>21</sub> with P103 | rBPI <sub>21</sub> with P104 | rBPI <sub>21</sub> with P105 | rBPI <sub>21</sub> with P123 |
| Oxacillin         | CSMHB       | 32 <sup>R</sup>                                       | 32 <sup>R</sup>             | <0.25 <sup>S</sup>           | 0.5 <sup>S</sup>             | 1 <sup>S</sup>               | 0.5 <sup>S</sup>             |
|                   | Serum       | 32 <sup>R</sup>                                       | > 32 <sup>R</sup>           | 32 <sup>R</sup>              | 32 <sup>R</sup>              | 32 <sup>R</sup>              | 32 <sup>R</sup>              |

- 52 -

Table 15

Effect of BPI protein product formulation on antibiotic susceptibility of *E. faecium*

|                   |             | Minimum Inhibitory Concentration (μg/mL) |                             |                              |                              |                              |                              |
|-------------------|-------------|--|-----------------------------|------------------------------|------------------------------|------------------------------|------------------------------|
| Antibiotic Tested | Medium Used | Control (no BPI)                         | rBPI <sub>21</sub> with F68 | rBPI <sub>21</sub> with P103 | rBPI <sub>21</sub> with P104 | rBPI <sub>21</sub> with P105 | rBPI <sub>21</sub> with P123 |
| Rifampicin        | CSMHB       | 4 <sup>R</sup>                           | 0.5 <sup>S</sup>            | 0.5 <sup>S</sup>             | 0.5 <sup>S</sup>             | 0.5 <sup>S</sup>             | 0.5 <sup>S</sup>             |
|                   | Serum       | 4 <sup>R</sup>                           | 1 <sup>S</sup>              | 1 <sup>S</sup>               | >4 <sup>R</sup>              | 0.5 <sup>S</sup>             | 0.5 <sup>S</sup>             |
| Chloramphenicol   | CSMHB       | 16 <sup>I</sup>                          | <4 <sup>S</sup>             | <4 <sup>S</sup>              | <4 <sup>S</sup>              | <4 <sup>S</sup>              | <4 <sup>S</sup>              |
|                   | Serum       | 8 <sup>S</sup>                           | 8 <sup>S</sup>              | 8 <sup>S</sup>               | 8 <sup>S</sup>               | 8 <sup>S</sup>               | 8 <sup>S</sup>               |
| Ciprofloxacin     | CSMHB       | 2 <sup>I</sup>                           | 1 <sup>S</sup>              | <0.5 <sup>S</sup>            | 1 <sup>S</sup>               | 1 <sup>S</sup>               | 1 <sup>S</sup>               |
|                   | Serum       | 2 <sup>I</sup>                           | 1 <sup>S</sup>              | 2 <sup>I</sup>               | 2 <sup>I</sup>               | 2 <sup>I</sup>               | 2 <sup>I</sup>               |

In additional experiments, a BPI protein product, rBPI<sub>21</sub>, was formulated with an anti-bacterial activity enhancing poloxamer, specifically PLURONIC P123, and with various concentrations of EDTA, and was evaluated for its effect on the antibiotic susceptibility of a *Pseudomonas*  
5 *aeruginosa* (ATCC 19660). Antibiotic susceptibility was determined using Microscan® panel plates (Dade Microscan, West Sacramento, CA) that allow simultaneous determination of minimum inhibitory concentrations for a number of different antibiotics.

The antimicrobial susceptibility tests performed on the  
10 Microscan® panel plates are miniaturizations of the broth dilution susceptibility test. Antimicrobial agents are serially diluted in Mueller-Hinton broth (supplemented with calcium and magnesium, or with sodium chloride for oxacillin, or with thymidine phosphorylase for trimethoprim, sulfamethoxazole and trimethoprim/ sulfamethoxazole) to concentrations bridging the range of  
15 clinical interest. One well on the 96-well Microscan® plate is a growth control well that contains dehydrated broth only. The remaining wells contain dehydrated broth and antibiotic (or broth and biochemical reagent indicator), which is rehydrated to the desired concentration by inoculation of a standardized suspension of test organism. The chromogenic biochemical agent  
20 indicators are used to identify and characterize the species of bacteria based on detection of pH changes and substrate utilization. After incubation overnight, the minimum inhibitory concentration (MIC) of an antibiotic for the test organism is determined by observing the well with the lowest concentration of the antibiotic that shows inhibition of growth. Gram-negative and gram  
25 positive organisms may be tested using any of the Microscan® panel plates (Microscan®, Dade Microscan, West Sacramento, CA). In these experiments with *P. aeruginosa*, the MIC Plus Type 2 panel plates were used. The concentrations of antibiotics tested in this panel plate are shown below in Table 16. The antibiotic susceptibility standards (interpretation of an MIC as  
30 resistant, intermediate or susceptible according to Microscan®'s NCCLS-

09696709-102400

derived standards) applicable to the gram-negative organisms that may be tested in each panel plate appear below in Table 16A.

09696709 " 102400

**Table 16**  
**ANTIBIOTIC CONCENTRATIONS TESTED IN**  
**MIC PLUS TYPE 2 PANEL PLATE**

|    | <b>Antibiotic</b>         | <b>Two-Fold Serial Dilutions Tested (μg/ml)</b> |
|----|---------------------------|---|
| 5  | Amoxicillin/K Clavulanate | 1/0.5-32/16                                     |
|    | Ampicillin/Sulbactam      | 1/0.5-32/16                                     |
|    | Azlocillin                | 64  |
|    | Aztreonam                 | 1-32  |
|    | Carbenicillin             | 16-128  |
| 10 | Cefamandole               | 1-32  |
|    | Cefonicid                 | 2-16  |
|    | Cefoperazone              | 4-32  |
|    | Cefotaxime                | 2-64  |
|    | Cefotetan                 | 4-32  |
| 15 | Ceftazidime               | 1-32  |
|    | Ceftizoxime               | 2-32  |
|    | Ceftriaxone               | 2-64  |
|    | Chloramphenicol           | 2-16  |
|    | Ciprofloxacin             | 0.25-4  |
| 20 | Imipenem                  | 0.5-16  |
|    | Mezlocillin               | 16-128  |
|    | Netilmicin                | 2-16  |
|    | Ticarcillin               | 16-128  |
|    | Ticarcillin/K Clavulanate | 16-128  |

09696709 102400



**Table 16A**  
**MICROSCAN MIC PLUS TYPE 2 ANTIBIOTIC**  
**SUSCEPTIBILITY RANGES FOR GRAM-NEGATIVE BACTERIA**

|    | Antibiotic                             | MIC (µg/ml) |              |             |
|----|--|-------------|--------------|-------------|
|    |  | Resistant   | Intermediate | Susceptible |
| 5  | Amoxicillin/K Clavulanate              | ≥ 32/16     | 16/8         | ≤ 8/4       |
|    | Ampicillin/Sulbactam                   | ≥ 32/16     | 16/8         | ≤ 8/4       |
|    | Azlocillin <sup>P</sup>                | > 64        |              | ≤ 64        |
|    | Aztreonam                              | ≥ 32        | 16           | ≤ 8         |
|    | Carbenicillin <sup>E</sup>             | ≥ 64        | 32           | ≤ 16        |
| 10 | Carbenicillin <sup>P</sup>             | > 128       |              | ≤ 128       |
|    | Cefamandole                            | ≥ 32        | 16           | ≤ 8         |
|    | Cefonicid                              | > 16        | 16           | ≤ 8         |
|    | Cefoperazone                           | > 32        | 32           | ≤ 16        |
|    | Cefotaxime                             | ≥ 64        | 16-32        | ≤ 8         |
| 15 | Cefotetan                              | > 32        | 32           | ≤ 16        |
|    | Ceftazidime                            | ≥ 32        | 16           | ≤ 8         |
|    | Ceftizoxime                            | > 32        | 16-32        | ≤ 8         |
|    | Ceftriaxone                            | ≥ 64        | 16-32        | ≤ 8         |
|    | Chloramphenicol                        | > 16        | 16           | ≤ 8         |
| 20 | Ciprofloxacin                          | ≥ 4         | 2            | ≤ 1         |
|    | Imipenem                               | ≥ 16        | 8            | ≤ 4         |
|    | Mezlocillin <sup>E</sup>               | ≥ 128       | 32-64        | ≤ 16        |
|    | Mezlocillin <sup>P</sup>               | ≥ 128       |              | ≤ 64        |
|    | Netilmicin                             | > 16        | 16           | ≤ 8         |
| 25 | Ticarcillin <sup>E</sup>               | ≥ 128       | 32-64        | ≤ 16        |
|    | Ticarcillin <sup>P</sup>               | ≥ 128       |              | ≤ 64        |
|    | Ticarcillin/K Clavulanate <sup>E</sup> | ≥ 128       | 32-64        | ≤ 16        |

004201" 60796960

| <b>Table 16A</b><br><b>MICROSCAN MIC PLUS TYPE 2 ANTIBIOTIC</b><br><b>SUSCEPTIBILITY RANGES FOR GRAM-NEGATIVE BACTERIA</b> |                          |              |             |
|--|--------------------------|--------------|-------------|
| Antibiotic   | MIC ( $\mu\text{g/ml}$ ) |              |             |
|  | Resistant                | Intermediate | Susceptible |
| Ticarcillin/K Clavulanate <sup>P</sup>   | $\geq 128$               |              | $\leq 64$   |

<sup>E</sup> *Enterobacteriaceae* only

<sup>P</sup> *Pseudomonas* only

For these experiments with *P. aeruginosa*, the following procedure was performed: The organism was streaked onto TSA plates (Remel, Lenexa, KN) and incubated for 18-24 hours overnight. Well-isolated colonies from the plates were emulsified in 3 ml of sterile Inoculum Water (catalog no. B1015-2, MicroScan® system, Dade Microscan, West Sacramento, CA) to a final turbidity equivalent to 0.5 McFarland Barium Sulfate standard. This cell suspension was vortexed for 2 to 3 seconds and 100  $\mu\text{l}$  was transferred to glass tubes containing 25 ml of Inoculum Water with Pluronic-D (catalog no. B1015-7, MicroScan® system, Dade Microscan, West Sacramento, CA) (hereinafter "Pluronic Inoculum Water"), or 25 ml of Pluronic Inoculum Water into which rBPI<sub>21</sub> in 0.2% PLURONIC P123, 0.002% TWEEN 80, 5mM sodium citrate, 150 mM NaCl ("rBPI<sub>21</sub>/P123") had been diluted to 64  $\mu\text{g/ml}$  rBPI<sub>21</sub>.

The 25 ml of this inoculum containing rBPI<sub>21</sub> was mixed by inversion and poured into a tray. The inoculum was drawn up into a manual 96-well pipetting system (RENOK™ rehydrator-inoculator system, Dade Microscan, West Sacramento, CA) designed for use with the Microscan® panel plates, and 110  $\mu\text{l}$  of the inoculum was delivered to each well of a Microscan® MIC Plus Type 2 panel plate. When added to the wells, this inoculum achieves a final bacterial concentration of  $4 \times 10^5$  to  $7 \times 10^5$  CFU/ml. The

004207-60296960

panel plates were then incubated at 35°C for 15-24 hours and read visually for cell growth.

No growth was defined as a slight whiteness in the well or a clear broth. Growth appeared as turbidity which could take the form of a white haze throughout the well, a white button in the center of the well, or a fine granule growth throughout the well. All wells were read against a black indirectly lighted background. Visual results of the biochemical reactions were read into a database for bacterial identification. The MICs for each antibiotic tested were determined by identifying the lowest concentration of antibiotic which inhibited visible growth.

Table 17 below displays a summary of the results of the antibiotic screening panel. The antibiotic susceptibility standards, which are the interpretation of an MIC as resistant, intermediate or susceptible according to Microscan®'s NCCLS-derived standards, are indicated in Table 16 as superscripts R, I and S, respectively. These data show that EDTA further enhanced the anti-bacterial activity of the rBPI<sub>21</sub>/P123 formulation by reversing resistance of the tested *P. aeruginosa* strain to cefonicid, cefotetan, cefamandole, chloramphenicol, ampicillin/sulbactam, and amoxicillin/k clavulanate, and by increasing the susceptibility of the tested *P. aeruginosa* strain to ceftizoxime, cefotaxime, ceftriaxone, and aztreonam.

09696709.102400

| <b>TABLE 17</b><br><b>Effects Of rBPI<sub>21</sub>/P123 Formulation ± Antibiotics On</b><br><i>P. aeruginosa</i> (ATCC 19660) with<br>varying concentrations of EDTA |  |                  |                  |                  |                 |  |
|--|--|------------------|------------------|------------------|-----------------|--|
| Antibiotic Tested  | Minimum Inhibitory Concentration of Antibiotic (µg/mL) |                  |                  |                  |                 |  |
|  | Control (No BPI <sub>21</sub> )                        | With 0% EDTA     | With 0.01% EDTA  | With 0.05% EDTA  | With 0.1% EDTA  |  |
| Ceftizoxime  | 32 <sup>I</sup>  | 16 <sup>I</sup>  | <2 <sup>S</sup>  | 8 <sup>S</sup>   | <2 <sup>S</sup> |  |
| Ceftazidime  | 2 <sup>S</sup>   | <1 <sup>S</sup>  | <1 <sup>S</sup>  | <1 <sup>S</sup>  | <1 <sup>S</sup> |  |
| Cefotaxime   | 32 <sup>I</sup>  | 16 <sup>I</sup>  | 4 <sup>S</sup>   | <2 <sup>S</sup>  | <2 <sup>S</sup> |  |
| Ceftriaxone  | 16 <sup>I</sup>  | 4 <sup>S</sup>   | 8 <sup>S</sup>   | <2 <sup>S</sup>  | <2 <sup>S</sup> |  |
| Cefoperazone   | <4 <sup>S</sup>  | <4 <sup>S</sup>  | <4 <sup>S</sup>  | <4 <sup>S</sup>  | <4 <sup>S</sup> |  |
| Cefonidic  | >16 <sup>R</sup>                                       | >16 <sup>R</sup> | <2 <sup>S</sup>  | <2 <sup>S</sup>  | <2 <sup>S</sup> |  |
| Cefotetan  | >32 <sup>R</sup>                                       | >32 <sup>R</sup> | <4 <sup>S</sup>  | <4 <sup>S</sup>  | <4 <sup>S</sup> |  |
| Netilmicin   | 4 <sup>S</sup>   | <2 <sup>S</sup>  | <2 <sup>S</sup>  | <2 <sup>S</sup>  | <2 <sup>S</sup> |  |
| Cefamandole  | >32 <sup>R</sup>                                       | >32 <sup>R</sup> | >32 <sup>R</sup> | >32 <sup>R</sup> | <4 <sup>S</sup> |  |

**TABLE 17**  
Effects Of rBPI<sub>21</sub>/PI23 Formulation ± Antibiotics On  
*P. aeruginosa* (ATCC 19660) with  
varying concentrations of EDTA

| Antibiotic Tested  | Minimum Inhibitory Concentration of Antibiotic (µg/mL) |                     |                     |                     |                     |
|--------------------|--|---------------------|---------------------|---------------------|---------------------|
|                    | Control (No BPI <sub>21</sub> )                        | With 0% EDTA        | With 0.01% EDTA     | With 0.05% EDTA     | With 0.1% EDTA      |
| Chloramphenicol    | > 16 <sup>R</sup>                                      | 8 <sup>S</sup>      | 16 <sup>I</sup>     | < 2 <sup>S</sup>    | < 2 <sup>S</sup>    |
| Ticarcillin        | < 16 <sup>S</sup>                                      | < 16 <sup>S</sup>   | < 16 <sup>S</sup>   | < 16 <sup>S</sup>   | < 16 <sup>S</sup>   |
| Azlocillin         | < 64 <sup>S</sup>                                      | < 64 <sup>S</sup>   | < 64 <sup>S</sup>   | < 64 <sup>S</sup>   | < 64 <sup>S</sup>   |
| Imipenem           | 1 <sup>S</sup>   | 1 <sup>S</sup>      | < 0.5 <sup>S</sup>  | < 0.5 <sup>S</sup>  | < 0.5 <sup>S</sup>  |
| Amp/Sulbact        | > 32 <sup>R</sup>                                      | > 32 <sup>R</sup>   | < 1 <sup>S</sup>    | 16 <sup>I</sup>     | 16 <sup>I</sup>     |
| Aztreonam          | 4 <sup>S</sup>   | 4 <sup>S</sup>      | < 1 <sup>S</sup>    | 2 <sup>S</sup>      | < 1 <sup>S</sup>    |
| Amox/K Clavulanate | > 32 <sup>R</sup>                                      | > 32 <sup>R</sup>   | < 1 <sup>S</sup>    | 32 <sup>R</sup>     | < 1 <sup>S</sup>    |
| Ciprofloxacin      | < 0.25 <sup>S</sup>                                    | < 0.25 <sup>S</sup> | < 0.25 <sup>S</sup> | < 0.25 <sup>S</sup> | < 0.25 <sup>S</sup> |

| <b>TABLE 17</b><br><b>Effects Of rBPI<sub>21</sub>/P123 Formulation ± Antibiotics On</b><br><i>P. aeruginosa</i> (ATCC 19660) with<br>varying concentrations of EDTA |  |                   |                   |                   |                   |
|--|--|-------------------|-------------------|-------------------|-------------------|
| Antibiotic Tested  | Minimum Inhibitory Concentration of Antibiotic (µg/mL) |                   |                   |                   |                   |
|  | Control (No BPI <sub>21</sub> )                        | With 0% EDTA      | With 0.01% EDTA   | With 0.05% EDTA   | With 0.1% EDTA    |
| Ticar/K Clavulanate  | < 16 <sup>s</sup>                                      | < 16 <sup>s</sup> | < 16 <sup>s</sup> | < 16 <sup>s</sup> | < 16 <sup>s</sup> |
| Mezlocillin  | < 16 <sup>s</sup>                                      | < 16 <sup>s</sup> | < 16 <sup>s</sup> | < 16 <sup>s</sup> | < 16 <sup>s</sup> |
| Carbenicillin  | 32 <sup>l</sup>  | < 16 <sup>s</sup> | < 16 <sup>s</sup> | < 16 <sup>s</sup> | < 16 <sup>s</sup> |

# EXAMPLE 7

## ANTI-BACTERIAL ACTIVITY OF COMPOSITIONS CONTAINING BPI PROTEIN PRODUCT AND POLOXAMER 188 OR POLOXAMER 403 ON *PSEUDOMONAS* INFECTION IN A RABBIT CORNEAL ULCERATION MODEL

The anti-bacterial activity of therapeutic compositions comprising BPI protein products with a poloxamer surfactant was evaluated in the context of administration both prior to and after *Pseudomonas* infection in a corneal infection/ulceration rabbit model.

For these experiments, the infectious organism was a strain of *Pseudomonas aeruginosa* 19660 obtained from the American Type Culture Collection (ATCC, Rockville, MD). The freeze dried organism was resuspended in nutrient broth (Difco, Detroit, MI) and grown at 37°C with shaking for 18 hours. The culture was centrifuged following the incubation in order to harvest and wash the pellet. The washed organism was Gram stained in order to confirm purity of the culture. A second generation was cultured using the same techniques as described above. Second generation cell suspensions were diluted in nutrient broth and adjusted to an absorbance of 1.524 at 600 nm, a concentration of approximately  $6.55 \times 10^9$  CFU/ml. A final  $1.3 \times 10^6$  fold dilution in nutrient broth yielded 5000 CFU/mL or  $1.0 \times 10^2$  CFU/0.02 mL. Plate counts for CFU determinations were made by applying 100 µL of the diluted cell suspension to nutrient agar plates and incubating them for 24-48 hours at 37°C.

The animals used were New Zealand White rabbits, maintained in rigid accordance to both SERI guidelines and the ARVO Resolution on the Use of Animals in Research. A baseline examination of all eyes was conducted prior to injection in order to determine ocular health. All eyes presented with mild diffuse fluorescein staining, characteristically seen in the normal rabbit eye. The health of all eyes fell within normal limits. Rabbits weighing between 2.5 and 3.0 kg were anesthetized by intramuscular injection of 0.5-0.7 mL/kg rodent cocktail (100 mg/mL ketamine, 20 mg/mL xylazine, and 10

mg/mL acepromazine). One drop of proparacaine hydrochloride (0.5% Ophthaine, Bristol-Myers Squibb) was applied to the eye prior to injection. Twenty microliters of bacterial suspension ( $1 \times 10^2$  CFU) prepared as described above was injected into the central corneal stroma of a randomly assigned eye while the other eye remained naive. Injections, simulating perforation of the corneal epithelium, were performed using a 30-gauge 1/2-inch needle and a 100  $\mu$ L syringe.

For the first series of experiments, a 5-day dosing regimen of BPI protein product (test drug) was as follows: on Day 0 of the study, 40  $\mu$ L of test drug or vehicle control was delivered to the test eye at 2 hours (-2) and 1 hour (-1) prior to intrastromal bacterial injection (time 0), then at each of the following 10 hours (0 through +9 hrs) post-injection for a total of 12 doses (40  $\mu$ L/dose); on each of Days 1-4 of the study, 40  $\mu$ L of test drug or vehicle control was delivered to the test eye at each of 10 hours (given at the same time each day, e.g., 8am-5pm). For these experiments, to test the poloxamer 188-containing therapeutic composition, 5 animals were treated with rBPI<sub>21</sub> (2 mg/mL in 5 mM citrate, 150 mM NaCl, 0.2% poloxamer 188, 0.002% polysorbate 80) and 5 with buffered vehicle, and to test the poloxamer 103-containing therapeutic composition, 5 animals were treated with rBPI<sub>21</sub> (2 mg/mL in 5 mM citrate, 150 mM NaCl, 0.2% poloxamer 403, 0.002% polysorbate 80) and 5 animals with placebo (5 mM citrate, 150 mM NaCl, 0.2% poloxamer 403, 0.002% polysorbate 80).

Eye examinations were conducted two times each day for each 5-day study via slit lamp biomicroscopy to note clinical manifestations. Conjunctival hyperemia, chemosis and tearing, mucous discharge were graded. The grading scale for hyperemia was: 0 (none); 1 (mild); 2 (moderate); and 3 (severe). The scale for grading chemosis was: 0 (none); 1 (visible in slit lamp); 2 (moderate separation); and 3 (severe ballooning). The scale for grading mucous discharge was: 0 (none) 1 slight accumulation); 2 (thickened discharge); and 3 (discrete strands). Photophobia was recorded as present or



absent. Tearing was recorded as present or absent. The corneal ulcer, when present, was assessed with respect to height (mm), width (mm), and depth (% of corneal thickness). Neovascularization was graphed with respect to the affected corneal meridians. Photodocumentation was performed daily as symptoms progressed throughout the experimental procedure.

At the completion of the 5-day study period, all rabbits were sacrificed via a lethal dose of sodium pentobarbital (6 grs/mL). Corneas were harvested and fixed in half-strength Karnovsky's fixative. The corneas were processed for light microscopy using Gram stain to assay for the presence of microbial organisms and using hematoxylin and eosin to assay for cellular infiltrate.

Examinations were conducted at 4, 24, 28, 48, 52, 72, 76, and 96 hours after injection of *Pseudomonas*. The results of these examinations are reported in Table 18 for the therapeutic composition comprising rBPI<sub>21</sub> with poloxamer 403, which provided the most potent effects.

004201-60796960

Table 18

| Summary of Clinical Observations<br>for therapeutic composition containing rBPI <sub>21</sub> and poloxamer 403 |                    |       |                    |       |                    |       |                         |              |                                |                              |
|---|--------------------|-------|--------------------|-------|--------------------|-------|-------------------------|--------------|--------------------------------|------------------------------|
|   | Hyperemia*         |       | Chemosis*          |       | Mucous*            |       | Neovas-<br>cularization |              | Ulcer Size<br>(mm)             |                              |
| Examination   | rBPI <sub>21</sub> | Plbo. | rBPI <sub>21</sub> | Plbo. | rBPI <sub>21</sub> | Plbo. | rBPI <sub>21</sub>      | Plbo.        | rBPI <sub>21</sub>             | Plbo.                        |
| Exam 1<br>4 hours   | 1.2                | 1.0   | 0.2                | 0.3   | 0.5                | 0     | None                    | None         | 1 ulcer<br>2mm                 | 1.4                          |
| Exam 2<br>24 hours  | 0.9                | 1.6   | 0.2                | 1.0   | 0.3                | 0.5   | None                    | None         | 1 ulcer<br>6mm                 | 3.4                          |
| Exam 3<br>28 hours  | 0.6                | 1.7   | 0.2                | 1.1   | 0.6                | 1.3   | None                    | None         | 1 ulcer<br>7mm                 | 5.2                          |
| Exam 4<br>48 hours  | 0.6                | 2.4   | 0.2                | 1.3   | 0.4                | 2.1   | None                    | None         | 1 ulcer<br>12mm<br>1 melt      | 11.4<br>3 melt<br>1 thinning |
| Exam 5<br>52 hours  | 0.8                | 2.4   | 0.2                | 1.2   | 0.2                | 1.6   | None                    | Yes<br>(1/5) | 1 ulcer<br>12mm<br>1 melt      | 11.4<br>3 melt<br>1 thinning |
| Exam 6<br>72 hours  | 0.6                | 2.4   | 0                  | 0.2   | 0.2                | 1.0   | None                    | Yes<br>(1/5) | 1 ulcer<br>12mm<br>melt & thin | 11.4<br>4 melt<br>1 thinning |
| Exam 7<br>76 hours  | 0.6                | 2.4   | 0                  | 0.2   | 0.2                | 0.8   | None                    | Yes<br>(2/5) | 1 ulcer<br>12mm<br>melt & thin | 11.4<br>4 melt<br>3 thinning |
| Exam 8<br>96 hours  | 0.6                | 2.4   | 0                  | 0.2   | 0.2                | 0.8   | None                    | Yes<br>(2/5) | 1 ulcer<br>12mm<br>melt & thin | 11.4<br>4 melt<br>3 thinning |

\* Mean scores of clinical observations graded on a scale of 0 (none) to 3 (severe).

004201" 60295960

The results set out in Table 18 reveal that treatment of the eye prior to and after perforation injury and injection of *Pseudomonas* provided substantial benefits in terms of reduced hyperemia, chemosis and mucous formation, as well as reduction in incidence of neovascularization along with reduced incidence and severity of corneal ulceration. At four hours after *Pseudomonas* injection, fluorescein staining of the cornea in both treated and control animals revealed small areas of staining consistent with the injection (puncture) injury. At 28 hours after injection, the rBPI<sub>21</sub>/poloxamer 403 treated eye evidenced clear ocular surfaces and typically were free of evidence of hyperemia, chemosis and mucous discharge while the vehicle treated eyes showed clouding of the ocular surface resulting from corneal edema and infiltration of white cells. Iritis was conspicuous in the vehicle treated eyes at 28 hours after injection and fluorescein dye application typically revealed areas of devitalized epithelium; severe hyperemia and moderate to severe chemosis and mucous discharge were additionally noted. At 48 hours after injection, mild hyperemia was sometimes noted in the rBPI<sub>21</sub>/poloxamer 403 treated eyes but mucous discharge and chemosis were absent; the rBPI<sub>21</sub>/poloxamer 403 treated corneas were otherwise typically clear and healthy appearing, as evidenced by the application of fluorescein dye. Vehicle treated eyes at 48 hours post infection displayed severe hyperemia, chemosis and mucous discharge were present; some corneas displayed corneal melting and thinning along with an ulcerating area clouded as a result of edema, cellular infiltration and fibrin deposition. At 52 hours following injection, rBPI<sub>21</sub>/poloxamer 403 treated eyes exhibited clear and healthy corneas which resisted staining with fluorescein, indicating that the formulation is safe and non-toxic to the corneal epithelium. In vehicle treated eyes at 52 hours post infection, sloughing of corneal epithelium was evident and while chemosis was decreasing, hyperemia was severe. In these experiments, several vehicle treated eyes presented with neovascularization, with vessels

09696709-102400

growing inward toward the central cornea. This manifestation was not noted in any rBPI<sub>21</sub>/poloxamer 403 treated eye.

5 Pathohistological evaluation of the rBPI<sub>21</sub>/poloxamer 403 treated corneas stained with hematoxylin and eosin revealed healthy, intact corneal epithelium and stroma; the tissue was free of white cell infiltration. In contrast, evaluation of the vehicle treated corneas revealed absence of an epithelium and extensive infiltration of white cells into the corneal stroma.

10 Additional pathohistological evaluation of the rBPI<sub>21</sub>/poloxamer 403 treated corneas stained with toluidine blue also revealed healthy, intact corneal epithelium and stroma, and further revealed corneal tissue free of *Pseudomonas* organisms. In contrast, evaluation of the vehicle treated corneas revealed rod shaped *Pseudomonas* organisms in the tissue and the presence of white cells advancing toward the organisms in the tissue. These results indicate effective corneal penetration of the  
15 rBPI<sub>21</sub>/poloxamer 403 and effective sterilization of the tissue without neovascularization.

20 The rBPI<sub>21</sub>/poloxamer 403 therapeutic composition tested in these experiments achieved the most dramatic beneficial antimicrobial and anti-angiogenic effects when compared with those of the rBPI<sub>21</sub>/poloxamer 188 therapeutic composition tested in this severe *Pseudomonas* injury/infection rabbit model. Benefits in terms of suppression of neovascularization were noted for treatment with the rBPI<sub>21</sub>/poloxamer 188 composition and no significant effects in reduction of hyperemia, chemosis, mucous formation and tearing were noted. The contrast in efficacy of the  
25 BPI<sub>21</sub>/poloxamer 403 composition with the lesser efficacy of the rBPI<sub>21</sub>/poloxamer 188 composition in these experiments suggested that formulation components, dosage and dosage regimen may all have a significant role in optimizing beneficial effects associated with methods according to the invention.

004201-60296960

### EXAMPLE 8

#### BACTERIAL AND FUNGAL GROWTH-INHIBITORY ACTIVITY OF COMPOSITIONS CONTAINING BPI PROTEIN PRODUCT AND POLOXAMER 188 OR POLOXAMER 403 IN THE PRESENCE OR ABSENCE OF EDTA

The antimicrobial preservative effectiveness of therapeutic compositions comprising BPI protein product and poloxamer surfactant were evaluated according to the U.S. Pharmacopeia (USP) microbiological test protocol (USP 23, [51] Antimicrobial Preservatives-Effectiveness, p. 1681) against the standard bacterial and fungal test microorganisms: *Escherichia coli* (ATCC No. 8739), *Pseudomonas aeruginosa* (ATCC No. 9027), *Staphylococcus aureus* (ATCC No. 6538), *Candida albicans* (ATCC No. 10231) and *Aspergillus niger* (ATCC No. 16404).

For these experiments, a small volume of the cultures from each of the five test microorganisms prepared according to the USP protocol was added into sterile containers with a solution of 2 mg/ml rBPI<sub>21</sub>, 0.2% poloxamer 188 (PLURONIC F68) or poloxamer 403 (PLURONIC P123), 0.002% TWEEN 80, 5mM sodium citrate and 150 mM sodium chloride. In some experiments, these solutions additionally contained various concentrations of EDTA. Aliquots of test solution were removed from the containers at various time periods after inoculation with the microorganisms (i.e., 7, 14, 21, and 28 days) and plated to determine the number of colony forming units (CFU) of each of the five microorganisms. According to USP standards, the product shows effectiveness if (a) the concentrations of viable bacteria are reduced to not more than 0.1% of the initial concentrations by the fourteenth day; (b) the concentrations of viable fungi remain at or below the initial concentrations during the first 14 days; and (c) the concentration of each test microorganism remains at or below these designated levels during the remainder of the 28-day test period.

09696709-102400

The results of initial testing of rBPI<sub>21</sub>/poloxamer 188 and rBPI<sub>21</sub>/poloxamer 403 compositions are shown in Tables 19A-19B below.

Table 19A

| CFUs after incubation with 2 mg/mL rBPI <sub>21</sub> /0.2% poloxamer 188 |                        |                        |                        |                        |                        |
|---|------------------------|------------------------|------------------------|------------------------|------------------------|
| Organisms   | Initial                | 7 Day                  | 14 Day                 | 21 Day                 | 28 Day                 |
| <i>E. coli</i>  | 4.9 x 10 <sup>6</sup>  | 1.67 x 10 <sup>3</sup> | 6.7 x 10 <sup>2</sup>  | <1                     | <1                     |
| <i>P. aeruginosa</i>  | 1.46 x 10 <sup>6</sup> | 1.7 x 10 <sup>2</sup>  | 5.8 x 10 <sup>3</sup>  | 4.7 x 10 <sup>4</sup>  | 2.05 x 10 <sup>5</sup> |
| <i>S. aureus</i>  | 3.6 x 10 <sup>6</sup>  | 7.5 x 10 <sup>2</sup>  | 7.8 x 10 <sup>1</sup>  | 2.9 x 10 <sup>2</sup>  | 1.15 x 10 <sup>3</sup> |
| <i>C. albicans</i>  | 3.3 x 10 <sup>6</sup>  | 2.62 x 10 <sup>6</sup> | 2.62 x 10 <sup>6</sup> | 2.96 x 10 <sup>6</sup> | 4.1 x 10 <sup>6</sup>  |
| <i>A. niger</i>   | 5.5 x 10 <sup>5</sup>  | 8.5 x 10 <sup>5</sup>  | 6.9 x 10 <sup>5</sup>  | 2.6 x 10 <sup>5</sup>  | 7.1 x 10 <sup>5</sup>  |

Table 19B

| CFUs after incubation with 2 mg/mL rBPI <sub>21</sub> /0.2% poloxamer 403 |                        |                       |                       |                       |                       |
|---|------------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| Organisms   | Initial                | 7 Day                 | 14 Day                | 21 Day                | 28 Day                |
| <i>E. coli</i>  | 7.2 x 10 <sup>5</sup>  | 0                     | 0                     | 0                     | 0                     |
| <i>P. aeruginosa</i>  | 1.02 x 10 <sup>5</sup> | 0                     | 0                     | 0                     | 0                     |
| <i>S. aureus</i>  | 6.2 x 10 <sup>5</sup>  | 1.8x 10 <sup>1</sup>  | 0                     | 0                     | 0                     |
| <i>C. albicans</i>  | 3.4 x 10 <sup>5</sup>  | 1 x 10 <sup>5</sup>   | 7.4 x 10 <sup>4</sup> | 7.9 x 10 <sup>4</sup> | 7.9 x 10 <sup>4</sup> |
| <i>A. niger</i>   | 1.9 x 10 <sup>5</sup>  | 1.5 x 10 <sup>5</sup> | 1.4 x 10 <sup>5</sup> | 1.4 x 10 <sup>5</sup> | 8.9 x 10 <sup>4</sup> |

When additional compositions of rBPI<sub>21</sub>/poloxamer 403 as described above were prepared with concentrations of 0.01%, 0.05% and 0.1% EDTA and tested in the experiments shown in Table 19B above, the results obtained were comparable to those shown in Table 19B above for all organisms.

In additional experiments, other compositions of 2mg/mL rBPI<sub>21</sub>, 0.2% PLURONIC P123, 0.002% TWEEN 80, 5mM sodium citrate, 150 mM sodium chloride with and without 0.05% EDTA were evaluated for effectiveness as described above. The results are shown in Table 20 below. In these experiments, 0.05% EDTA further enhanced the antimicrobial effectiveness of the rBPI<sub>21</sub>/poloxamer 403 composition.

- 70 -

TABLE 20

| CFUs after incubation with 2 mg/mL rBPI <sub>21</sub> /0.2% poloxamer 403 ± 0.05% EDTA |                        |                       |                        |                        |                        |
|--|------------------------|-----------------------|------------------------|------------------------|------------------------|
| Organisms  | Initial                | 7 Day<br>-EDTA/+ EDTA | 14 Day<br>-EDTA/+ EDTA | 21 Day<br>-EDTA/+ EDTA | 28 Day<br>-EDTA/+ EDTA |
| <i>E. coli</i>   | 1.97 x 10 <sup>5</sup> | 10                    | 1                      | 1                      | 1                      |
| <i>P. aeruginosa</i>   | 7 x 10 <sup>4</sup>    | 1                     | 1                      | 1                      | 1                      |
| <i>S. aureus</i>   | 9.4 x 10 <sup>4</sup>  | 1                     | 3.9 x 10 <sup>3</sup>  | 1                      | 1                      |
| <i>C. albicans</i>   | 3 x 10 <sup>3</sup>    | 8.8 x 10 <sup>3</sup> | 1.5 x 10 <sup>3</sup>  | 4.1 x 10 <sup>4</sup>  | 5 x 10 <sup>5</sup>    |
| <i>A. niger</i>  | 7.25 x 10              | 1.8 x 10 <sup>4</sup> | 1.2 x 10 <sup>4</sup>  | 1.69 x 10 <sup>4</sup> | 1.4 x 10 <sup>4</sup>  |
|  |                        |                       | 1.45 x 10 <sup>4</sup> | 3.4 x 10 <sup>2</sup>  | 1.7 x 10 <sup>2</sup>  |
|  |                        |                       | 5.7 x 10 <sup>4</sup>  | 4.4 x 10 <sup>4</sup>  | 1.66 x 10 <sup>4</sup> |

Numerous modifications and variations of the above-described invention are expected to occur to those of skill in the art. Accordingly, only such limitations as appear in the appended claims should be placed thereon.

09696709 102400